

The **16th** Annual Advanced

# Pain Conference

and **Practical Workshop**

29-31 August, 2011, **Budapest, Hungary**


[www.congressline.hu/pain2011](http://www.congressline.hu/pain2011)

Program Book  
and Syllabus



The **19th**  
**WIP FIPP**  
**Examination**

1 September, 2011  
Budapest, Hungary



## CONTENT

Greetings .....	4.
WIP Council .....	6.
Faculty .....	7.
Acknowledgement .....	8.
Exhibitor & Sponsor Profiles .....	9.
General Information .....	11.

### Detailed program

29 August .....	14.
30 August .....	15.
31 August .....	16.
Awards Ceremony .....	17.
Syllabus .....	21.
Authors Index .....	53.



# GREETINGS

## Dear Colleagues and Friends,

Ten years ago we were ready to have our 6th Annual Budapest Conference when the news of 9/11 changed the world through terror that was so unnecessary. It seems the world has lost a great deal of innocence. The first thought on that Tuesday morning was that we had to cancel the conference scheduled for September 16-19, 2001. I was already in Budapest in the organizing days of the conference. But, under pressure from the news, I felt the decision could not be made that day and I would make a decision in two days. The following morning, on Wednesday, the world looked so bad that I would have been ready to cancel except that I had made the decision to wait two days. On that second day, Thursday, September 13, I woke up more determined that no terrorist will prevent us from our mission and sent the word out that we are not canceling our meeting. It was truly amazing that the lecturers came through sheer determination and difficulties of travel, not to be prevented from our mission of becoming better physicians. We came from all over Europe and North America, South America, the Middle East and Far East to participate. The participation of our German, British and Spanish colleagues was heart-warming love. Because of their drive, the individuals who have been to Budapest have gotten better and are today making major impact in their local societies, resulting in making patients better.

The fact is clear that physicians who are better trained in the technical procedures are providing better patient care around the world. Over the years we have not only learned how to do procedures but more importantly how to teach them. This year's tenth anniversary since 9/11 makes us even more determined to become better. I am grateful to every one of you who have become Fellows of Interventional Pain Practice (FIPP). We now have 630 FIPPs worldwide. I especially congratulate the Hungarian section for meeting the required numbers to be a Section. Our local arrangement chairperson, Dr. Edit Racz, has been relentless in her determination to find x-ray machines for us when we needed them, also for going annually to the Ethics Committee in Budapest to represent the Interventional Pain Conference and get permission for us to do our practice sessions. We are grateful for the opportunity that has been awarded us during these years and especially grateful for the permission of Semmelweis University Medical School and the cooperation of all of faculty and staff for use of their facilities.

The leadership of WIP Past President, Dr. Serdar Erdine, MD, FIPP, has been remarkable in bringing more open organizational structure to WIP. The spiritual father of WIP, Prithvi Raj, MD, FIPP, is forever present and it is no small achievement of his to have come up with full tax-exempt approval of the World Institute of Pain Foundation as a charitable organization. We already have our physicians channeling individual patient contributions to the Foundation so that we may help others. The future of the WIP Foundation will bring notable and unimaginable support to deserving people around the world.

The Budapest Conference has always focused on practical and clinically-relevant material. We are interested in safety and efficacy. New and effective pain treatment modalities will be brought to the participants in a setting of morning lectures and afternoon practice of the procedures under expert guidance.

The WIP World Congress this year had to be cancelled in Seoul, Korea because of travelling concerns for so many would-be participants. The Congress is rescheduled to February 4-6, 2012 in Miami Beach, Florida. Please remember this date, a unique opportunity to meet old friends and make new ones, as well as keep abreast of the very latest in our field of study.

Our WIP journal Pain Practice is moving forward with greater and better acceptance and relevance. The Journal will publish the awards for Centers of Excellence. Keep in mind to get your center positioned for this highly coveted award.

The Budapest Conference organizational committee is working hard to make this an outstanding conference again, and we are going to make sure that 2011 will be another memorable experience and conference. Budapest is one of the most beautiful cities in the world. The city is changing for the better. The air is cleaner. The food is outstanding, and the music is fitting for the occasion for you to be a participant in our annual reunion in Budapest.

I have to thank our sponsors that make this annual educational and learning opportunity possible. Our mission simply would not have been possible without the loyal and continuing support of our sponsors. We could not have come this far in such few years. I ask you to note those who sponsor us and make every effort to them.

Please remember that we are not competing with other groups or societies as we all belong to multiple societies. We exist purely because there is a need for activities. Our patients need us to learn in order to improve care we provide them. The WIP is firmly committed to those who wish to teach and those who wish to learn so we can help our patients. Pain knows no boundaries. People in every country need pain doctors who can help them and make the quality of life better worldwide.

In Hungary, we shall watch and listen to the most beautiful starry-eyed children of the Hungarian Folk Dance Group who will show us how dancing should be done. You will be better, and you will make us better by your participation, and your patients will know the difference because your ability will improve through participation. Welcome to Budapest.

**Gabor B. Racz, MD, FIPP**

Director of Budapest Conference  
Past President World Institute of Pain  
Co-Director Texas Tech University Health  
Sciences Center, Lubbock, Texas

**Ricardo Ruiz-López, MD, FIPP**

President World Institute of Pain  
Director Tramiento del Dolor  
Clinica del Dolor de Barcelona

# WIP COUNCIL

## President WIP

**Ricardo Ruiz-López**, MD, FIPP

## Executive Board

**Ricardo Ruiz-López**, MD, FIPP, Founder and President – Spain

**Serdar Erdine**, MD, FIPP, Founder and Immediate Past President – Turkey

**Gabor B. Racz**, MD, FIPP, Founder and Past President – USA

**P. Prithvi Raj**, MD, FIPP, Founder and Past President – USA

**Charles A. Gauci**, MD, FRCA, FIPP, FFPMRCA, Honorary Secretary – UK

**Kris C. P. Vissers**, MD, PhD, FIPP, Honorary Treasurer – The Netherlands

**Craig T. Hartrick**, MD, DABPM, FIPP, Editor-in-Chief, Pain Practice – USA

**Miles R. Day**, MD, FIPP, DABA, Chair, Board of Examination – USA

**Ira B. Fox**, MD, FIPP, Chair, Advisory Board – USA

**José R. Rodríguez Hernández**, MD, FIPP, Chair, Board of Sections – Puerto Rico

**Dianne Willard**, Executive Officer – USA

## Section Chairmen

**Meir Bennun**, MD, FIPP – Israel

**Frantz J. Cólimon**, MD, FIPP – Colombia

**Peter G. Courtney**, MBBS, FIPP – Australia

**Gautam Das**, MD, FIPP – India

**Neels de Villiers**, MD, FRCA, FIPP – Africa

**Magdi Ramzi Iskander**, MD, FFARCS, FIPP – Middle East

**Sang Chul Lee**, MD, PhD, FIPP – Korea-Japan-China

**Martin Marianowicz**, MD, FIPP – Central-Eastern Europe

**Philippe Mavrocordatos**, MD, FIPP – Switzerland

**Patrick R. McGowan**, MBChB, FRCA, FIPP, FFPMRCA – UK

**Charles Amaral de Oliveira**, MD, FIPP – Latin America

**Nuri Süleyman Özyalçın**, MD, FIPP – Turkey

**Mahdi Panah Khahi**, MD, FIPP – Iran

**Edvin Koshi**, MD, FIPP – Canada

**Teresa Bovaira**, MD, FIPP – Spain

**Edit Racz**, MD, FIPP – Hungary

**José R. Rodríguez Hernández**, MD, FIPP – Puerto Rico

**Andrea M. Trescot**, MD, FIPP – USA

**Athina Vadalouca**, MD, PhD, FIPP – Mediterranean

**Jan Van Zundert**, MD, PhD, FIPP – Benelux

**Alex Sow Nam Yeo**, MD, PhD, FIPP – SE Asia

## WIP Examination Board

**Chair: Miles R. Day**, MD, FIPP, DABA

**Co-Chair: Maarten van Kleef**, MD, PhD, FIPP

**Liaison to WIP: Serdar Erdine**, MD, FIPP

**James E. Heavner**, DVM, PhD, FIPP (Hon)

**Charles Amaral de Oliveira**, MD, FIPP

**Neels de Villiers**, MD, FIPP

**Sang Chul Lee**, MD, PhD, FIPP

**Patrick R. McGowan**, MBChB, FRCA, FIPP, FFPMRCA

**Jan van Zundert**, MD, PhD, FIPP

## Congress Organizers

**Program Director: Gabor B. Racz**, MD, FIPP

**Co-Director: James E. Heavner**, DVM, PhD, FIPP (Hon)

## Local Arrangement Committee

**Chair: Edit Racz**, MD, FIPP

**Agnes Stogicza**, MD, FIPP

**Lorand Eross**, MD, PhD, FIPP

## FACULTY

**Mert Akbas**, MD, FIPP

**Adnan A. Al-Kaisey**, MB ChB, FFRCA, FPMRCA, FIPP

**Ray M. Baker**, MD, FIPP

**Hemmo A. Bosscher**, MD, FIPP

**Cosimo Bruni**, MD

**Kenneth B. Chapman**, MD, FIPP

**Serdar Erdine**, MD, FIPP

**Lorand Eross**, MD, PhD, FIPP

**Joseph D. Fortin**, MD

**James E. Heavner**, DVM, PhD, FIPP (HON)

**Rafael Justiz**, MD, MS, FIPP, DABIPP

**Sang Chul Lee**, MD, PhD, FIPP

**Liong Liem**, MD, FIPP

**John Nelson**, MD, FIPP

**Edit Racz**, MD, FIPP

**Gabor B. Racz**, MD, FIPP

**Richard Rauck**, MD, FIPP

**Olav J. J. M. Rohof**, MD, PhD, FIPP

**Murray H. Rosenthal**, DO, FAPA

**Ricardo Ruiz-López**, MD, FIPP

**Andrea Trescot**, MD, FIPP

**Kris C. P. Vissers**, MD, PhD, FIPP

**Jan Peter Warnke**, MD

# ACKNOWLEDGEMENTS

The 16th Annual Advanced Interventional Pain Conference and Practical Workshop gratefully acknowledge the following companies for their support of this event:



**Boston  
Scientific**



**CoMEDICAL** **COSMAN** MEDICAL  
Radiofrequency Pain Management Radiofrequency Pain Management



**Epimed**  
*Designed by Experts for Experts*



**Kimberly-Clark**



**Medtronic**



**ML** MILLENNIUM  
LABORATORIES  
*because pain matters*



**ST. JUDE MEDICAL**  
MORE CONTROL. LESS RISK.



**ziehm imaging**  
dedicated to clinical innovation

# EXHIBITORS AND SPONSORS PROFILES

## **Boston Scientific**

**25155 Rye Canyon Loop, Valencia, CA 91355, USA**

[www.controlyourpain.com](http://www.controlyourpain.com)

Boston Scientific's Precision Plus™ SCS System powered by SmoothWave™ Technology blends sophistication and simplicity to deliver life-changing therapy for chronic pain patients. Investing in innovative products, clinical initiatives, and world-class service, Boston Scientific is committed to Making life smoother™ for physicians, patients, and the Neuromodulation community.

## **CoMedical – Cosman Medical**

**Gieterijstraat 46, Ridderkerk 2984 AB, Netherlands**

[www.comedical.eu](http://www.comedical.eu), [www.cosmanmedical.com](http://www.cosmanmedical.com)

CoMedical and Cosman Service Centre is based in Ridderkerk nearby Rotterdam and is a specialized medical company that focuses on development and distribution of Radiofrequency equipment as well specific catheters and needles for the minimal invasive treatment of Chronic Pain and publishing books for pain management. We have an interactive relationship with anesthesiologists, neurosurgeons, pain managers, research and industry. We are exclusive distributor for Cosman Medical, Oakworks and Epimed in the Benelux. Now also are the publisher of the 3rd Edition of Manual of RF Techniques of Dr. Charels A. Gauci MD. FRCA. FIPP. FPPMRCA. Since the eighties Enrico Cohen founder and CEO of CoMedical is involved in minimal invasive surgery and specialized on pain management. Since many years he cooperate with Cosman medical (Radionics) now he is also Vice President Sales for Cosman Medical in EAME. CoMedical Team can provide products, training and service for you, your staff and equipment.

## **Epimed International Inc.**

**Crossroads Business Park 141 Sal Landrio Drive Johnstown, NY 12095, USA**

[www.epimedpain.com](http://www.epimedpain.com)

Epimed International, inc will be featuring products designed for chronic and acute pain management techniques. We will display the Expanded Line of Racz® Spring Guide Epidural Catheters; RX™, R.K.™, and FIC Epidural Introducer Needles; R-F™ Line of Radiofrequency Products; Coudé™ & Straight Blunt Nerve Block Needles; and Mini Trays. Also being shown are Radiation Safety Products and Anatomical Models.

## **Kimberly-Clark Health Care**

**1400 Holcomb Bridge Road, Roswell, Georgia 30076, USA**

[www.rfpainmanagement.com](http://www.rfpainmanagement.com)

KIMBERLY-CLARK\* Pain Management offers a complete line of chronic pain solutions, including an innovative radiofrequency system that features standard, pulsed, and multi-level RF lesioning functions. Our revolutionary cooled-RF technology is designed to treat symptomatic patients experiencing discogenic, sacroiliac, thoracic, and lumbar z-joint pain. Kimberly-Clark's pain management probes are reusable, durable, and steam-sterilizable. Our radiopaque RF cannulae provide enhanced visualization during positioning. At Kimberly-Clark Health Care, we deliver innovative healthcare solutions that healthcare professionals depend on to meet the demands of a fast-paced world. Our solutions help prevent, diagnose, and manage major issues in Infection Prevention, Digestive Health and Pain Management.

## Medtronic International Trading Sarl

Route de Molliau 31, 1131 Tolochenaz, Switzerland

[www.medtronic.com](http://www.medtronic.com)

We're committed to Innovating for life by pushing the boundaries of medical technology and changing the way the world treats chronic disease. To do that, we're thinking beyond products and beyond the status quo - to continually find more ways to help people live better, longer.

## Millennium Laboratories Inc.

16981 Via Tazon, San Diego, CA 92127, USA

[www.milleniumpulaboratories.com](http://www.milleniumpulaboratories.com)

Millennium Laboratories, Inc. is research-based medication monitoring company whose test panels, technology, educational resources and scientific experts are specifically focused on clinicians who treat chronic pain. Millennium is the only major Urine Drug Testing (UDT) laboratory exclusively utilizing the latest cutting-edge technology (LC-MS/MS) allowing for the fastest turn-around time in the industry.

## St. Jude Medical

[sjmneuro.com](http://sjmneuro.com)

St. Jude Medical Neuromodulation Division develops implantable systems that put more control into the hands of those who treat chronic pain patients. We offer a comprehensive suite of neurostimulation products for the management of chronic intractable pain, exceptional service, technical support, and professional education programs.

## Ziehm Imaging GmbH.

Donaustrasse 31, 90451 Nuernberg, Germany

[www.ziehm.com](http://www.ziehm.com)

Ziehm Imaging specializes in the development, manufacturing and worldwide marketing of mobile X-ray-based imaging solutions. The company has been market leader in Germany for many years as well as in many other European countries. Today, Ziehm Imaging is a global systems provider, employing over 280 people worldwide. Extensive in-house development know-how is reflected in the Ziehm Imaging C-arms' high medical imaging performance, intelligent generator technology, significant dose savings and seamless digital network integration. Building on competence and creativity, as well as continuous dialog and close cooperation with renowned universities, research centers and hospitals, Ziehm Imaging has developed groundbreaking technologies that have made the company a global trendsetter in intelligent interventional imaging. Please visit [www.ziehm.com](http://www.ziehm.com) for more information.

# GENERAL INFORMATION

## Dates & Venue

**The 16th Annual Advanced Interventional Pain Conference & Practical Workshop**

29 – 31 August, 2011

## Conference Site

**Kempinski Hotel Corvinus Budapest – Regina Ball Room**

H-1051 Budapest, Erzsébet tér 7-8.

## Practical Workshop

**University Labs**

H-1091 Budapest, Üllői út 93.

Daily bus transfers are provided within the venues.

## The 19th WIP FIPP Examination

1 September, 2011

**Venue: University Labs**

H-1091 Budapest, Üllői út 93.

## Conference Website

[www.congressline.hu/pain2011](http://www.congressline.hu/pain2011)

## Language

English

## CME Credits

18 credits were granted by the European Accreditation Council for Continuing Education (EACCME).

## Opening Hours of the Registration Desk at Hotel Kempinski

Sunday, 28 August	14.00 – 19.30
Monday, 29 August	07.00 – 13.30
Tuesday, 30 August	07.00 – 13.30
Wednesday, 31 August	07.00 – 13.30

## FIPP Exam Registration at Kempinski Hotel

Wednesday, 31 August	16.00-19.00
----------------------	-------------

## Registration Fee

(Regular Fees after 15 July, 2011)

Pain Conference & Practical Workshop	1600 Euro
Pain Conference	1150 Euro
Accompanying person fee	350 Euro
FIPP Exam registration fee	2500 USD

## Internet

Free of charge Wi-Fi service available at the venue.

## Commercial Exhibition

The exhibition will be opened from Monday, 29 August until 31 August at the Kempinski Hotel foyer. Delegates will have the opportunity to meet representatives of pharmaceutical and diagnostic equipment companies at their stands to discuss new developments and receive up-to-date product information.

## Meals

Coffee breaks, lunches, welcome cocktail and award ceremony dinner are included in the registration fee.

## Hotels

**Kempinski Hotel Covinus Budapest\*\*\*\*\*** (Conference venue)

H-1051 Budapest, Erzsébet tér 7-8.

[www.kempinski-budapest.com](http://www.kempinski-budapest.com)

**Hotel Central Basilica\*\*\***

H-1051 Budapest, Hercegprímás u. 8.

[www.hotelcentral-basilica.hu](http://www.hotelcentral-basilica.hu)

## Official Social Events

### Faculty Dinner

**Sunday**, 28 August, 2011, 20.00-22.00 (only for Faculty Members)

**Robinson Restaurant**

**Dress Code:** business casual

**Meeting point:** Kempinski Hotel lobby at 18.30 (Bus transportation is provided.)

### Welcome Cocktail

**Monday**, 29 August, 2011, 20.00-22.00 (for all registered guests)

**Kempinski Hotel, Regina Ball Room**

**Programme:** Csillagszemű Dance Ensemble

**Dress Code:** Business casual

### Awards Ceremony Dinner

**Tuesday**, 30 August, 2011, 20.00-23.00 (for all registered guests)

**Hungarian Academy of Sciences Ball Room** (1014 Budapest, Országház u. 28-30.)

**Programme:** Award Ceremony and String Quartet

**Dress Code:** formal

**Meeting point:** Kempinski Hotel lobby at 19.30 (Bus transportation is provided.)

### Sightseeing Tour in Budapest

**Monday**, 29 August, 2011, 09.30-13.00

**Price: 30 Euro / person** (Minimum number of participants: 15 persons)

Including in the accompanying person fee

An approximately 4-hour long sightseeing tour, which shows the most attractive features of the capital.

Transportation by bus, with English speaking guide, refreshment and all entrance fees are included.

**Meeting point:** Kempinski Hotel lobby at 09.15 (conference venue)

## Useful Information

### How to get to the Conference Venue?

To reach the Conference Venue there are several means of transport:

**Metro station "Deák Ferenc Square"** station of Metro line 1,2,3.

**From the airport to the conference venue use the Airport Minibus Service**, fixed rates for passengers

One way: 2990 HUF = cca 12 Euro, Return ticket: 4990 HUF = cca 18 Euro

Tel: +36 1 296 8555 [www.airportshuttle.hu](http://www.airportshuttle.hu) or use the PAIN2011 Official Taxi Company:

**City Taxi** +36 1- 211-1111 (Rate: **5100-5300HUF = cca 23-25Euro**).

### Climate

The climate of Budapest is continental. In August usually nice warm weather can be expected with a max. temperature of 25-30 °C, while the lowest temperature during the night ranging between 15-20 °C.

Nevertheless some rainy days can be expected.

### Insurance

The registration fees do not include provision for the insurance of participants against personal accidents, illness, cancellation, theft, property loss or damage. Participants are advised to take adequate personal travel insurance.

### Currency

The Forint (HUF), the official national currency, is convertible. The exchange rates applied in Budapest banks, official exchange offices and hotels may vary. All the major credit cards are accepted in Hungary in places displaying the emblem at the entrance.

Exchange rate: 1 Euro = 270 HUF in August, 2011.

### Credit Cards

In general, VISA, EC/MC and American Express credit cards are accepted in most restaurants, cafés, shops and petrol stations.

### Stores and Shopping

The opening hours of Budapest stores are generally 10.00-18.00 on weekdays and 10.00-13.00 on Saturday. The shopping centers are open from 10.00-21.00 from Monday to Saturday and from 10.00-18.00 on Sunday.

### Electricity

The voltage in Hungary is 230V, 50 Hz AC.

### Parking

If you drive a personal or rented car, always try to park at a guarded parking lot and do not leave any valuables in the car. Please note, that Budapest is divided into paying areas, with one parking meter in each street. The maximum parking time duration is 2 hours, tariffs may vary.

# Detailed Program

MONDAY, 29 August, 2011

Regina Ball Room

- 07:40**      **Opening Remarks**  
**Gabor B. Racz**, MD, FIPP, Program Director  
**Ricardo Ruiz-López**, MD, FIPP, President of WIP  
**Edit Racz**, MD, FIPP, Chair Local Arrangement Committee
- Moderator:**    **Lorand Eross**, MD, PhD, FIPP
- 07:50**      **Future of Interventional Pain Practice**  
**Richard Rauck**, MD, FIPP
- 08:30**      **Percutaneous interventions for trigeminal neuralgia**  
**Serdar Erdine**, MD, FIPP
- 09:00**      **Intraspinal Drug Delivery**  
**Richard Rauck**, MD, FIPP
- 09:30**      **Discography**  
**Olav J. J. M. Rohof**, MD, PhD, FIPP
- 10:00**      **Minimally Invasive Techniques for Discogenic Pain**  
**Ricardo Ruiz-López**, MD, FIPP
- 10:30**      **Coffee Break**
- 11:00**      **Vertebral Body Stabilization Techniques**  
**Rafael Justiz**, MD, MS, FIPP, DABIPP
- 11:30**      **Lysis of Epidural Adhesions: How to Do It and How to Teach it Safely**  
**Gabor B. Racz**, MD, FIPP
- 12:00**      **RF of Joints and Discs**  
**Olav J. J. M. Rohof**, MD, PhD, FIPP
- 12:30**      **Sacroiliac Pain**  
**Joseph D. Fortin**, MD
- 13:00**      **Lunch**
- 13:30**      **Transport to University**  
**Labs Afternoon workshops**

# Detailed Program

TUESDAY, 30 August, 2011

Regina Ball Room

- Moderator:**    **Mert Akbas**, MD, FIPP
- 08:00**      **Complications of Interventional Pain Therapy**  
**John Nelson**, MD, FIPP
- 08:30**      **Epidural Injections**  
**Kris C. P. Vissers**, MD, PhD, FIPP
- 09:00**      **Imaging for Interventional Pain Management**  
**Joseph D. Fortin**, MD
- 09:30**      **Controversies in the Diagnosis of Painful Lumbar Disc Degeneration**  
**Ray M. Baker**, MD, FIPP
- 10:00**      **Joint Pain and Its Treatment with Biological Agents and Drugs**  
**Cosimo Bruni**, MD
- 10:30**      **Coffee Break**
- 11:00**      **Lumbosacral Spinal Endoscopy Round Table**  
Moderator: **Gabor B. Racz**, MD, FIPP  
Science Background – **James E. Heavner**, DVM, PHD, FIPP (HON)  
Clinical Perspective – **Hemmo A. Bosscher**, MD, FIPP  
Application to Arachnoiditis – **Jan Peter Warnke**, MD
- 12:00**      **Cold Allodynia, Nerve Entrapment Radiculopathies and Illusions**  
The Science – **James E. Heavner**, DVM, PhD, FIPP (HON)  
The Clinical – **Gabor B. Racz**, MD, FIPP
- 12:30**      **Lunch**
- 13:30**      **Transport to University**  
**Labs Afternoon workshops**

# Detailed Program

WEDNESDAY, 31 August, 2011

Regina Ball Room

- Moderator:** Kenneth B. Chapman, MD, FIPP
- 08:00**      **Management of Spinal Stenosis**  
John Nelson, MD, FIPP
- 08:30**      **Pathophysiology and Pharmacologic Management of Neuropathic Pain**  
Andrea Trescot, MD, FIPP
- 09:00**      **Pharmacologic Management of Cancer Pain**  
Kris C. P. Vissers, MD, PhD, FIPP
- 09:30**      **Current Neurosurgical Approaches to the Treatment of Pain**  
Lorand Eross, MD, PhD, FIPP
- 10:00**      **Facial Pain and Cervicogenic Headache**  
Liong Liem, MD, FIPP
- 10:30**      **Coffee Break**
- 11:00**      **Ultrasound Guided Treatment**  
Sang Chul Lee, MD, PhD, FIPP
- 11:30**      **Sacral Nerve Root Stimulation – Recent Advances**  
Gabor B. Racz, MD, FIPP
- 12:00**      **High Frequency Spinal Cord Stimulation in the Management of Axial Back Pain**  
Adnan A. Al-Kaisey, MB ChB, FFRCA, FPMRCA, FIPP
- 12:30**      **Urine Drug Screening**  
Murray H. Rosenthal, DO, FAPA
- 13:00**      **Lunch**
- 13:30**      **Transport to University**  
Labs Afternoon workshops

# FIPP Awards Ceremony

TUESDAY, 30 August, 2011, at 20:00

**Master of Ceremonies:** Miles Day, MD, FIPP, DABA  
**Opening Remarks – Local Organizing Committee**  
Edit Racz, MD, FIPP  
Agnes Stogicza, MD, FIPP  
Lorand Eross, MD PhD, FIPP

**The First Ten Years of FIPP**  
P. Prithvi Raj, MD, FIP, WIP Founder & Past President

**Presentation of Certificates to Fellows of Interventional Pain Practice (FIPP)**  
WIP Board of Examination Members

## FIPP Class of 2 September, 2010 – Budapest, Hungary

- 602 Aertssen, Jack**, MD, FIPP      The Netherlands
- 603 Bhatia, Anuj**, MBBS DNB MD FRCA MNAMS FPPMRCA, FIPP      Canada
- 604 Blau, Jonathan**, MD, FIPP      USA
- 605 Cada, Miroslav**, MD, FIPP      Austria
- 606 Canty, Tim**, MD, FIPP      USA
- 607 Deshmukh, Prakash**, MD, FIPP      India
- 608 Dirckx, Veerle**, MD, FIPP      Belgium
- 609 Du Plessis, Pauline**, MD, FIPP      South Africa
- 610 Filippini-de Moor, Gertruda**, MD, FIPP      The Netherlands
- 611 Gautam, Sujeet Kumar Singh**, MD, FIPP      India
- 612 Goswami, Subrata**, MBBS, FIPP      India
- 613 Haghshenas, Seyed Mohammad**, MBBS, FIPP      Iran
- 614 Hussain, Assad**, MBBS, FHKCA, FANZCA, Dip Pain Mgt (HKCA), FPPMANZCA, FIPP      Iran
- 615 Kal, Jasper Emiel**, MD, PhD, FIPP      The Netherlands
- 616 Kanthed, Pravesh**, MBBS, DA, DNB, FIPP      India
- 617 Laverdière, Mélanie**, MD, FIPP      Switzerland
- 618 Lim, Kyung-Joon**, MD, PhD, FIPP      South Korea

<b>619 Mestrum, Roel</b> , HM, MD, FIPP	Belgium
<b>620 Mishriqy, Basem Henry Habib</b> , MD, FIPP	Egypt
<b>621 Naderi -Nabi, Bahram</b> , MD, FIPP	Iran
<b>622 Nimmakayalu Sampu</b> , Swaroop, FCARCSI FIPP	UK
<b>623 Perumal, Muralitharan</b> , MD, FIPP	Malaysia
<b>624 Phua, Darren Shingkuan</b> , MD, FIPP	Canada
<b>625 Ray, Subrata</b> , DA, FIPP	India
<b>626 Reddy, Raja Varadapura</b> , MD, FRCA, FPPMRCA, FIPP	UK
<b>627 Shanthanna, Harsha</b> , MD, DNB, FIPP	Canada
<b>628 Sie, Robert Bernhard</b> , MD, FIPP	The Netherlands
<b>629 Soin, Amol</b> , MD, FIPP	USA
<b>630 Tay, Teik Guan</b> , MBBS, FIPP	Australia

## FIPP Class of 25 June, 2011 – Maastricht, The Netherlands

<b>631 Annehmohammadzadeh, Haligaldy</b> , MD, FIPP	Iran
<b>632 Balthasar, Andrea Johanna</b> , MD, FIPP	Germany
<b>633 Dakheel, Raad</b> , MD, FIPP	Belgium
<b>634 Husada, Satria</b> , MD, FIPP	The Netherlands
<b>635 Janssen, Markus</b> , MD, FIPP	The Netherlands
<b>636 Joseph, A. Alister</b> , FCARCSI, FIPP	Ireland
<b>637 Kallewaard, Jan-Willem</b> , MD, FIPP	The Netherlands
<b>638 Köder, Brigitte</b> , MD, FIPP	The Netherlands
<b>639 Lim, Tai Chuan</b> , MD, FIPP	The Netherlands
<b>640 Mehta, Palak Bhavin</b> , MD, FIPP	India
<b>641 Meyer, Johannes</b> , MD, FIPP	South Africa
<b>642 Munshi, Shirazahmed Mo.Abbas</b> , MD, FIPP	India
<b>643 Oberholzer, Andries Johannes</b> , MD, FIPP	South Africa
<b>644 Patel, Anant Gunvantbhai</b> , MD, FIPP	India
<b>645 Perez-Martinez, Jordi</b> , MD, PhD, FIPP	UK

<b>646 Rouwet, Edward</b> , MD, FIPP	The Netherlands
<b>647 Sahinler, Bolkar E.</b> , MD, FIPP	USA
<b>648 Dongen, Vincent Ran</b> , MD, FIPP	The Netherlands
<b>649 Wilson, Eric</b> , MB, BCh, FIPP	South Africa

Big things  
come in small  
packages.

With its all-in-one design, Ziehm Solo is one of the most compact and versatile C-arms for crowded treatment scenarios.

Equipped with state-of-the-art imaging technology, this C-arm delivers great performance for orthopedics, traumatology, pain management and other applications. Versatile viewing options enable you to configure the unit according to your needs and local conditions. All this at a competitive price.

→ For more information visit [www.ziehm.com](http://www.ziehm.com)



## Syllabus

## SERDAR ERDINE, MD, FIPP

### BIOGRAPHICAL SKETCH

Prof. Serdar Erdine MD,FIPP is the immediate past president of WIP and CEO of the World Institute of Pain Foundation. He is the chairman of the Istanbul Pain Center affiliated with Istanbul Bilim University.

### PERCUTANEOUS INTERVENTIONS FOR TRIGEMINAL NEURALGIA

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- What sets percutaneous interventions for trigeminal neuralgia apart from other interventional pain therapies
- The rationale for performing percutaneous interventions for trigeminal neuralgia
- Indications for each percutaneous intervention for trigeminal neuralgia
- Expected outcomes of percutaneous interventions for trigeminal neuralgia
- Complications of percutaneous interventions for trigeminal neuralgia
- State of the art for percutaneous interventions for trigeminal neuralgia

#### Key Points

Definition of a good result is not easy

- Variation of the follow-up also affects the decision making
- Learning curve-experience is important
- No method is curable
- Cure is unpredictable, recurrence rate is predictable
- All techniques have advantages and disadvantages
- RF is still the preferred technique
- It is less morbid more cost effective than open techniques
- Glycerol causes mild complications, less effective
- Balloon causes mild sensorial loss
- Recurrence rate for open techniques may be similar

#### References

1. Erdine S. Targets and optimal imaging for cervical spine and head blocks. *Techniques in Regional Anesthesia and Pain Management*. 2007, 11: 63-72.
2. Erdine S. Sympathetic blocks of the head and neck. In: Raj PP et al. (eds) *Interventional pain management, image guided procedures*. Philadelphia, Saunders-Elsevier. 2008, 108-127.
3. Erdine S, Racz GB, Noe CE. Somatic blocks of the head and neck. In: Raj PP et al. (eds) *Interventional Pain Management, Image Guided Procedures*. Philadelphia, Saunders-Elsevier. 2008, 77-108.
4. Hakanson S. Trigeminal neuralgia treated by the injection of glycerol into the trigeminal cistern. *Neurosurgery*. 1981, 9: 638-646.
5. Kanpolat Y, Savas A, Bekar A, et al. Percutaneous controlled radiofrequency trigeminal rhizotomy for the treatment of idiopathic trigeminal neuralgia: 25-year experience with 1,600 patients. *Neurosurgery*. 2001, 48: 524-532; discussion 532-534.
6. Latchaw JP Jr, Hardy RWJ, Forsythe SB, et al. Trigeminal neuralgia treated by radiofrequency coagulation. *J Neurosurg*. 1983, 59: 479-484.
7. Lopez BC, Hamlyn PJ, Zakrzewska JM. Systemic review of ablative neurosurgical techniques for the treatment of trigeminal neuralgia. *Neurosurgery*. 2004, 54: 973-983.
8. Narouze S. Complications of head and neck procedures. *Techniques in Regional Anesthesia and Pain Management*. 2007, 1: 171-177.
9. Ruiz-López R, Erdine S. Treatment of cranio-facial pain with radiofrequency procedures. *Pain Practice*. 2002, 2: 206-213.
10. Sweet WH, Wepsic SG. Controlled thermocoagulation of trigeminal ganglion and rootlets for differential destruction of pain fibers. Part 1: Trigeminal neuralgia. *J Neurosurg*. 1977, 40: 143-156
11. Van Kleef M, Van Genderen WE, Narouze S, Nurmikko TJ, Van Zundert J, Geurts JW, Mekhail N. Trigeminal Neuralgia. *Pain Practice*. 2009, 9: 252-259

## OLAV J. J. M. ROHOF, MD, PHD, FIPP

### BIOGRAPHICAL SKETCH

Dr. Olav Rohof (04-06-1950) MD, PhD, FIPP is an anesthesiologist and interventional pain specialist, and head of the pain clinic of Orbis Medical Center in Sittard Geleen, the Netherlands.

As an anesthesiologist he has been working full time in the pain clinic since 1989.

He has trained many colleagues from all over the world.

He has special interest and experience in trigeminal neuralgia and facial pain, back pain, neck pain and headache, treatment of joints and CRPS. In 2002 he wrote his thesis on the RF treatment of the ganglion Gasseri in trigeminal neuralgia (University of Bonn Germany).

### DISCOGRAPHY

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

1. Definitions: Disc stimulation, Discography, provocative discography, disc manometry(1)
2. Patient selection, in- and exclusion criteria
3. Procedure and technique
4. The level(s) to be tested?
5. In disc stimulation use of disc manometry is advised, if available
6. Evaluation of results of disc stimulation and discography
7. Complications
8. Differential Diagnosis
9. Discography still indicated?
10. Results of a prospective pilot study of pulsed RF disc treatment in the nucleus following lumbar discography

#### Key points:

- Definitions
  - **Disc stimulation:** a procedure developed to define a painful intervertebral disc by intradiscal injection .
  - **Discography:** injection of dye to describe the internal disc morphology.
  - **Provocative Discography:** a combination of both of the above.
  - **Disc manometry :** disc stimulation with intradiscal pressure measurement.
  - Only to be used when interventional treatment for reduction of discogenic pain is considered.
- Patient selection: chronic low back pain with or without (pseudo)radicular irradiation for longer than 3 months, not reacting on conservative therapy and medication, in which facet- and SI-joint pain have been excluded or treated without sufficient result.
- Procedure and technique will be presented.
- Combination of history, physical examination and additional examinations (MRI, EMG, CT) determines the probably involved levels to be tested.
- A suspected normal level functions as control level.
- Disc manometry, technique and criteria will be discussed (2,3)
- Evaluation of results of disc stimulation will be discussed according to ISIS and IASP criteria.
- Evaluation of discography using the Dallas Discogram Scale (5)
- Discitis is the mostly feared complication, prophylactic use of antibiotics is advised in international guidelines, intravenously or intradiscally, in the "two needle technique" questionable (4)

- DD: trauma, fracture, tumour, infection, neurologic disease, visceral pain
- Despite controversial literature lumbar (provocative) discography and disc manometry still is the gold standard for confirming the diagnosis discogenic pain. A strong patient selection is needed to improve the results of invasive intradiscal treatments.
- Technique and results of a prospective pilot study of Pulsed RF disc treatment in the nucleus following lumbar discography

#### References

1. J.van Zundert ,F.Huygen,J.Patijn, M.van Kleef: Praktische richtlijnen anesthesiologische pijnbestrijding, gebaseerd op klinische diagnoses; Lumbosacraal-Discogene lage ruggpijn ,JW Kallewaard, M.Terheggen, G.J.Groen,M.Sluijter, M.van Kleef..Pijn Kennis Centrum Maasticht 2009
2. Seo KS, Derby R, Date ES, Lee SH, Kim BJ, Lee CH. In vitro measurement of pressure differences using manometry at various injection speeds during discography. *Spine J.* 2007; 7: 68-73.
3. O'Neill C, Kurgansky M. Subgroups of positive discs on discography. *Spine.* 2004; 29: 2134-9.
4. Willems PC, Jacobs W, Duinkerke ES, De Kleuver M. Lumbar discography: should we use prophylactic antibiotics? A study of 435 consecutive discograms and a systematic review of the literature. *J Spinal Disord Tech.* 2004; 17: 243-7.
5. Sachs BL, Vanharanta H, Spivey MA, Guyer RD, Videman T, Rashbaum RF, Johnson RG, Hochschuler SH, Mooney V. Dallas discogram description. A new classification of CT/discography in lowback disorders. *Spine.* 1987; 12: 287-94.

#### Further relevant literature:

6. Kuslich SD, Ulstrom CL, Michael CJ. The tissue origin of low back pain and sciatica: a report of pain response to tissue stimulation during operations on the lumbar spine using local anesthesia. *Orthop Clin North Am.* 1991; 22: 181-7.
7. Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The relative contributions of the disc and zygapophyseal joint in chronic low back pain. *Spine.* 1994; 19: 801-6.
8. Cohen SP, Larkin TM, Barna SA, Palmer WE, Hecht AC, Stojanovic MP. Lumbar discography: a comprehensive review of outcome studies, diagnostic accuracy, and principles. *Reg Anesth Pain Med.* 2005; 30: 163-83.
9. Groen GJ, Baljet B, Drukker J. Nerves and nerve plexuses of the human vertebral column. *Am J Anat.* 1990; 188, (3): 282-96.
10. Oh WS, Shim JC. A randomized controlled trial of radiofrequency denervation of the ramus communicans nerve for chronic discogenic low back pain. *Clin J Pain.* 2004; 20: 55-60.
11. Nakamura SI, Takahashi K, Takahashi Y, Yamagata M, Moriya H. The afferent pathways of discogenic low-back pain. Evaluation of L2 spinal nerve infiltration. *J Bone Joint Surg Br.* 1996; 78: 606-12.
12. Ohnmeiss DD, Vanharanta H, Ekholm J. Relation between pain location and disc pathology: a study of pain drawings and CT/discography. *Clin J Pain.* 1999; 15: 210-7.
13. Ohnmeiss DD, Vanharanta H, Ekholm J. Relationship of pain drawings to invasive tests assessing intervertebral disc pathology. *Eur Spine J.* 1999; 8: 126-31.
14. Vanharanta H. The intervertebral disc: a biologically active tissue challenging therapy. *Ann Med.* 1994; 26: 395-9.
15. Zhou Y, Abdi S. Diagnosis and minimally invasive treatment of lumbar discogenic pain--a review of the literature. *Clin J Pain.* 2006; 22: 468-81.
16. Carragee EJ, Hannibal M. Diagnostic evaluation of low back pain. *Orthop Clin North Am.* 2004; 35: 7-16.
17. Ito M, Incorvaia KM, Yu SF, Fredrickson BE, Yuan HA, Rosenbaum AE. Predictive signs of discogenic lumbar pain on magnetic resonance imaging with discography correlation. *Spine.* 1998; 23: 1252-8; discussion 9-60.
18. Gilbert FJ, Grant AM, Gillan MG, Vale LD, Campbell MK, Scott NW, Knight DJ, Wardlaw D. Low back pain: influence of early MR imaging or CT on treatment and outcome--multicenter randomized trial. *Radiology.* 2004; 231: 343-51.
19. Wolfer LR, Derby R, Lee JE, Lee SH. Systematic Review of Lumbar Provocation Discography in Asymptomatic Subjects with a Metaanalysis of False-positive Rates. *Pain Physician.* 2008; 11:513-38.
20. O'Neill C, Kurgansky M. Subgroups of positive discs on discography. *Spine.* 2004; 29: 2134-9.

Additional literature to be posted on the website

## RICARDO RUIZ-LÓPEZ, MD, FIPP

### BIOGRAPHICAL SKETCH

Ricardo Ruiz-López, MD, Neurosurg., FIPP, is Director of Barcelona Spine and Pain Institute (Institut de Columna Vertebral / Clínica del Dolor de Barcelona), Executive Member of the Board of Directors of Hospital Delfos (Barcelona) and CEO Project for Barcelona Spine & Pain Surgery Clinic.

After receiving his MD degree from the University of Madrid in 1975 and the Board of Neurosurgery in 1980, he founded in 1986 Clínica del Dolor de Barcelona.

His major areas of scientific interest are the Neurosurgery of Pain, the Interventional Techniques and Surgery for Spinal Chronic Pain Conditions, and the development of new organizational models for Patient 's Care.

Editor of a number of medical journals, he has published extensively on Pain Management and Interventional Pain Therapies.

He is a Founding Member of various National and International Medical Societies on the Pain Field, and Visiting Professor and Lecturer at European and American Universities.

President of the Organizing Committee of the II EFIC Congress (European Federation of IASP Chapters) "Pain in Europe" Barcelona, September 1997 and of the 3rd World Congress on Pain of WIP (World Institute of Pain), Barcelona, September 2004.

President of World Institute of Pain (WIP) 2011-2014, President of the Catalan Pain Society (Catalonia, Spain) 2006-2010, and Permanent Trustee of the World Institute of Pain Foundation, NC. USA.

### MINIMALLY INVASIVE TECHNIQUES FOR DISCOGENIC PAIN

Minimally invasive spine surgery has evolved rapidly within the last two decades in an effort to decrease morbidities associated with open procedures.

Low back pain of discogenic origin is a highly prevalent condition. Internal disc disruption has been described as the internal disruption of disc architecture of the disc without signs of disc protrusions or without positive signs for nerve root compression. Tears that appear in the layers of the annulus fibrosus allow nuclear material to irritate sensitive sinu-vertebral nerve-endings. Concentric annular tears and rim lesions are often present. The modified Dallas discogram classification is now the gold standard for the computer tomography (CT) classification of annular tears. It classifies discal lesions into five possible severities of the radial annular tear. Provocation discography is the most accurate diagnostic procedure, with two components to it: an attempt to reproduce the patient 's pain by injecting contrast material into the disc, and to perform a painless discogram in the adjacent disc. Gadolinium-enhanced magnetic resonance imaging is an alternative to CT discogram.

There are several therapeutic options for discogenic pain, ranging from conservative management to open surgery. In this review we described most percutaneous techniques that are currently being used to treat discogenic pain with and without disc herniation, as well as some of the emerging minimally invasive surgery techniques for disc removal.

#### References

1. Knoeller SM, Seifried C. History of spine surgery. *Spine* 25:2838-2843, 2000.
2. Fang HSY, Ong GB. Direct anterior approach of the upper cervical spine. *J Bone Joint Surg Am.* 44\*:1588-1604, 1962.
3. Roh SW, Kim DH, Cardoso AC, et al. Endoscopic foraminotomy using MED system in cadaveric specimens. *Spine* 25:260-264, 2000.
4. Gogan WJ, Fraser RD, Chymopapain: a 10-year, double-blind study. *Spine* 27:722-731, 1992.
5. Van der Belt H, Franssen S, Deutman R. Repeat chemonucleolysis is safe and effective. *Clin Orthop* 363:121-125, 1999.

6. Ascher PW, Heppner F. CO2 laser in neurosurgery. *Neurosurg Rev* 7:123-133, 1984.
7. Crock HV. A reappraisal of intervertebral disc lesions. *Med J Aust* 1:983-989, 1970.
8. Crock HV. Internal disc disruption: A challenge to disc prolapse fifty years on. *Spine* 11:650-653, 1986.
9. Schwarzer AC, Aprill CN, Derby R, et al. The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine* 20:1878-1888, 1995.
10. Merskey H, Bogduk N. Classification of Chronic Pain: Descriptions of Chronic Pain Síndromes and definitions of Pain Terms. Seattle: IASP Press, 1994.
11. Ruiz-López R, Pichot C. Percutaneous Therapeutic Procedures for Disc Lesions. In: PP Raj, et al, eds. *Interventional Pain Management: Image-Guided Procedures*. Saunders, 2nd ed. 2008. Pp 539-558.
12. Lee KS, Doh JW, Bae HG, et al. Diagnostic criteria for the clinical síndrome of IDD. *Br J Neurosurg* 17:19-23, 2003.
13. Osti OL, Vernon-Roberts B, Fraser RD. Volvo-Award-anulus tears and intervertebral disc degeneration: an animal model. *Spine* 15:762-766, 1990.
14. Moore RJ, Osti OL, Vernon-Roberts. Osteoarthritis of the facet joint resulting from anular rim lesions. *Spine* 24:519-524, 1999.
15. Moore RJ, et al. Remodeling of vertebral bone alter outer anular injury in sheep. *Spine* 21:936-940, 1996.
16. Key JA, Ford LT. Experimental intervertebral disc lesions. *J Bone Joint Surg* 30a:621, 1984.
17. Moore RJ, Osti OL, Vernon-Roberts B, et al. Changes in endplate vascularity alter an outer anulus tear in sheep. *Spine* 17:874-877, 1992.
18. Kim KS, Yoon ST, Li J, et al. Disc degeneration in the rabbit: a biochemical and radiological comparison between two disc injury models. *Spine* 30:33-37, 2005.
19. Sachs BL, et al. Dallas discogram description: a new classification of CT/discography in low back disorders. *Spine* 12:287-294, 1987.
20. Aprill C, Bogduk N. High intensity zone. *Br J Radiol* 65:361-369, 1992.

Additional references to be posted on the website.

## RAFAEL JUSTIZ, MD, MS, FIPP, DABIPP

### BIOGRAPHICAL SKETCH

Dr. Rafael Justiz is currently the Director of Interventional Pain Management , Department of Neurosciences, Saint Anthony's Hospital, Oklahoma City, Oklahoma.

Dr Justiz earned a Bachelor and Masters in Sciences from Florida International University in Miami, Florida, and his Doctor of Medicine from Medical college of Wisconsin. He completed his anesthesia residency at the University of South Florida in Tampa, and received his fellowship in Interventional Pain Management at Texas Tech University in Lubbock, Texas. Dr. Justiz joined the faculty at the international pain institute at University Health Sciences Center and now is currently in private practice.

He is board-certified in anesthesiology by the American Board of Anesthesiology and has Added Qualifications in Pain Management by the same board. He also holds the WIP Fellow in Interventional Pain Practice certification (FIPP) and is a Diplomate of the American Board of Interventional Pain Physicians (ABIPP).

Dr Justiz has published several book chapters and journal articles. His areas of interest's include peripheral field/spinal cord stimulation and treatment of refractory head and facial pain.

### VERTEBRAL BODY STABILIZATION TECHNIQUES

#### Objectives

Upon completion of this presentation attendees will be able to discuss

- Osteoporosis
- Treatment options for osteoporosis
- Vertebral Augmentation
- Identify patient and workup
- Different Techniques
- How to perform vertebral augmentation
- Complications

#### Key Points

- Discuss osteoporosis including risk factors, epidemiology, its economic effects and clinical consequences. Look at the guidelines for determining osteoporosis, and be able to recognize the disease process and what treatment options there are available.
- Discuss ideal patient selection and workup, and define fracture configurations.
- Discuss different imaging modalities that can be used and their differences.
- Discuss how vertebral augmentation reduces pain and what mechanism are involved.
- Look at the indications, contraindications and relative contraindications involved with vertebral augmentation.
- Discuss the different techniques employed in vertebral body augmentation, transpedicular and extrapedicular approaches. Look at the anatomical landmarks and proper imaging technique for safety. In detail define how each technique is performed and the approaches that can be employed including proper trajectory and vertebral access.
- Recognize the common complications and practice safe techniques to avoid these complication

#### References

1. NIH. Osteoporosis prevention, diagnosis, and therapy. NIH Consensus Statement. 2000;17:1-45.
2. USPST. Screening for osteoporosis: U.S. preventive services task force recommendation statement. *Ann Intern Med*. 2011;154:356-64. Epub 2011 Jan 17.
3. Lewiecki EM, Bilezikian JP, Khosla S, Marcus R, McClung MR, Miller PD, Watts NB, Maricic M. Osteoporosis update from the 2010 Santa fe bone symposium. *J Clin Densitom*. 2011;14:1-21.

4. Becker DJ, Kilgore ML, Morrisey MA. The societal burden of osteoporosis. *Curr Rheumatol Rep.* 2010;12:186-91.
5. Leslie WD, Schousboe JT. A Review of Osteoporosis Diagnosis and Treatment Options in New and Recently Updated Guidelines on Case Finding Around the World. *Curr Osteoporosis Rep.* 2011 Jun 8. [Epub ahead of print]
6. National Osteoporosis Foundation: Osteoporosis: What is it? Washington DC: National Osteoporosis Foundation. March 2004.
7. Kanis JA, Borgstrom F, De Laet C, Johansson H, Johnell O, Jonsson B, Oden A, Zethraeus N, Pflieger B, Khaltav N. Assessment of fracture risk. *Osteoporosis Int.* 2005;16:581-9. Epub 2004 Dec 23.
8. Blume SW, Curtis JR. Medical costs of osteoporosis in the elderly Medicare population. *Osteoporosis Int.* 2011;22:1835-44. Epub 2010 Dec 17.
9. Hargunani R, Le Corroller T, Khashoggi K, Murphy KJ, Munk PL. Percutaneous vertebral augmentation: the status of vertebroplasty and current controversies. *Semin Musculoskeletal Radiol.* 2011;15:117-24. Epub 2011 Apr 15.
10. Kim HS, Kim SH, Ju CI, Kim SW, Lee SM, Shin H. The role of bone cement augmentation in the treatment of chronic symptomatic osteoporotic compression fracture. *Korean Neurosurg Soc.* 2010;48:490-5. Epub 2010 Dec 31.
11. Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res.* 1993;8:1137-48.
12. Rölinghoff M, Zarghooni K, Schlüter-Brust K, Sobottke R, Schlegel U, Eysel P, Delank KS. Indications and contraindications for vertebroplasty and kyphoplasty. *Arch Orthop Trauma Surg.* 2010;130:765-74. Epub 2010 Mar 11.
13. McGraw JK, Cardella J, Barr JD, Mathis JM, Sanchez O, Schwartzberg MS, Swan TL, Sacks D; SIR Standards of Practice Committee. Society of Interventional Radiology quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol.* 2003;14:827-31.
14. Barbero S, Casorzo I, Durando M, Mattone G, Tappero C, Venturi C, Gandini G. Percutaneous vertebroplasty: the follow-up. *Radiol Med.* 2008;113:101-13. Epub 2008 Feb 25.
15. Monticelli F, Meyer HJ, Tutsch-Bauer E. Fatal pulmonary cement embolism following percutaneous vertebroplasty (PVP). *Forensic Sci Int.* 2005 20;149:35-8.
16. Amoretti N, Hovorka I, Marcy PY, Grimaud A, Brunner P, Bruneton JN. Aortic embolism of cement: a rare complication of lumbar percutaneous vertebroplasty. *Skeletal Radiol.* 2007;36:685-7. Epub 2007 Mar 30.
17. Deramond H, Saliou G, Aveillan M, Lehmann P, Vallée JN. Respective contributions of vertebroplasty and kyphoplasty to the management of osteoporotic vertebral fractures. *Joint Bone Spine.* 2006;73:610-3. Epub 2006 Oct 11.
18. Burton AW. Vertebroplasty and Kyphoplasty: Case Presentation, Complications, and Their Prevention. *Pain Medicine* 2008;9:S58-64.
19. Uppin AA, Hirsch JA, Centenera LV, Pfeifer BA, Pazianos AG, Choi IS. Occurrence of new vertebral body fracture after percutaneous vertebroplasty in patients with osteoporosis. *Radiology.* 2003;226:119-24.

## OLAV J. J. M. ROHOF, MD, PHD, FIPP

### (PULSED) RF TREATMENT OF DISC AND JOINTS

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

1. An overview of RF and PRF disc treatment techniques (1)
2. Treatment of articular pain in small and large joints (2)
3. Causes of articular pain, prevalence, pathophysiology
4. New technique of PRF treatment of the Lateral Atlanto Axial (LAA) joint
5. Results of 1 yr follow up of first 100 cases of PRF AA joint treatment (5) Technique PRF treatment Knee, shoulder, trapezio-metacarpal, and hallux valgus (2)
6. Results PRF treatment Knee, shoulder, trapezio-metacarpal, and hallux valgus (2)
7. Shoulder treatment with PRF suprascapular nerve, on indication combined with the one entry – 3 compartment block .

#### Key points

1. There is equivocal evidence about RF heating techniques of the disc, a prospective pilot study on PRF disc treatment shows promising results, but this should be confirmed in a randomised study.(6)
2. Osteoarthritis (OA) is the most common type of joint disease (22% of the adult population), pain is the major symptom determining functional loss; there is a poor correlation between pain and radiological changes; Infiltration with steroids are short lasting (3)
3. Pulsed RF treatment of small joints like the lateral AA joint without any injection is more accessible in the supine position with a true lateral fluoroscopic approach , and has remarkable relatively long-lasting results. (4, 5)
4. Diagnostic tools: History, clinical examination, radiology, blood tests and biomarkers.
5. Approaches for PRF treatment of the knee, shoulder, trapezio-metacarpal and metatarso-halyngeal joints will be presented.
6. Results of PRF treatment (2) will be presented.
7. Small joints respond particularly well to PRF treatment
8. Results of PRF treatment of Shoulder and Knee joints are superior to steroid infiltrations
9. Discussion: PRF treatment of large joints combined with viscosuppletion with hyaluronic acid, but without corticosteroids.

#### Literature

1. Raj P.P. Intervertebral disc: anatomy-physiology-pathophysiology-treatment. *Pain Pract.* 2008;8:18-44.
2. Schianchi Pietro M.: New frontiers of PRF: treatment of articular pain in small and large joints, International Symposium Invasive Procedures in Motion 2011, 21-22 January, Notwill Switzerland
3. Hepper C.T., Halvorson J.J. et al. The efficacy and duration of intra-articular corticosteroid injection for knee osteoarthritis: a systematic review of level I studies. *J.Am Acad Ortop Surg* 2009; 17:638-46.
4. Halim W., Chua NH, Vissers KC. Long-term pain relief in patients with cervicogenic headaches after pulsed radiofrequency application into the lateral atlantoaxial (C1-2) joint using an anterolateral approach. *Pain Pract.* 2010; 10:267-71.
5. Rohof OJJM . PRF treatment of the Lateral AA joint , one year follow up of first 100 cases with a true lateral approach. (personal communication, WIP London June 2009)
6. Rohof OJJM . Intradiscal pulsed radiofrequency application following provocative discography for the management of degenerative disc disease and concordant pain. *Pain Pract* 2011 (submitted).

Epiducer™  
Lead Delivery System



## INDUSTRY-FIRST LEAD DELIVERY SYSTEM FOR UNPRECEDENTED ACCESS TO LEAD OPTIONS.

St. Jude Medical's Epiducer lead delivery system is the next generation in epidural access. This revolutionary system allows introduction of a unique class of paddle leads as well as multiple lead arrays through a single percutaneous entry point. With unparalleled options and reduced procedural complexity, the Epiducer system is redefining percutaneous procedures.

[sjmneuro.com](http://sjmneuro.com)



Not all products approved in all countries. Please ask your local representative

**Indications for Use:** Chronic, intractable pain of the trunk and limbs. **Contraindications:** Demand-type cardiac pacemakers, patients who are unable to operate the system or who fail to receive effective pain relief during trial stimulation. **Warnings/Precautions:** Diathermy therapy, cardioverter defibrillators, magnetic resonance imaging (MRI), explosive or flammable gases, theft detectors and metal screening devices, lead movement, operation of machinery and equipment, postural changes, pediatric use, pregnancy, and case damage. Patients who are poor surgical risks, with multiple illnesses, or with active general infections should not be implanted. **Adverse Events:** Painful stimulation, loss of pain relief, surgical risks (e.g., paralysis). Clinician's manual must be reviewed prior to use for detailed disclosure. **Caution:** U.S. federal law restricts this device to sale and use by or on the order of a physician.

Epiducer is a trademark of Advanced Neuromodulation Systems, Inc. d/b/a St. Jude Medical Neuromodulation Division. ST. JUDE MEDICAL, the nine-squares symbol, and MORE CONTROL. LESS RISK. are registered and unregistered trademarks and service marks of St. Jude Medical, Inc. and its related companies. ©2011 St. Jude Medical, Inc. All rights reserved.

## JOSEPH D. FORTIN, MD

### BIOGRAPHICAL SKETCH

Dr. Fortin is Medical Director of Spine Technology and Rehabilitation and Clinical Professor of Indiana University School of Medicine, Fort Wayne, Indiana.

### SACROILIAC PAIN 2011

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- The unique anatomical and biomechanical features of the sacroiliac joint.
- Key clinical features of patient presenting with symptomatic sacroiliac joints.
- Diagnostic evaluation of SIJ dysfunction.
- Rehabilitation of SIJ dysfunction.
- Interventional techniques for addressing SIJ pain.

#### Key Points

- The sacroiliac joint is a putative source of low back pain and sciatica. The prevalence of sacroiliac joint dysfunction is between two and thirty percent.
- The SIJ is a true synovial joint with hyaline cartilage on the sacral side and fibrocartilage on the iliac side. The articular surfaces have numerous interdigitating ridges and depressions.
- The ventral ligaments are a thin extension of the capsule and the dorsal ligaments are a series of discontinuous bands.
- Mechanically, the sacroiliac joint is a relay station transmitting loads to and from the trunk and lower extremities.
- Histological analysis of the sacroiliac joint has verified the presence of nerve fibers within the joint capsule and joint ligaments. While the levels of innervation and divisions of SIJ innervation have been the subject of debate. A fetal correlate study found no receptors in the ventral capsule and concluded that the SIJ is innervated by the sacral dorsal rami.
- No physical exam findings are unequivocally pathognomonic of sacroiliac joint pain. The Fortin Finger Test is a reliable indicator of sacroiliac joint pain, yet it lacks sensitivity.
- Controlled diagnostic blocks appear to be the sole direct method of distinguishing symptomatic versus asymptomatic sacroiliac joints.
- Rehabilitative measures include manual medicine techniques, pelvic stabilization exercise to allow dynamic postural control, and muscle balancing of the trunk and lower extremities.
- Interventional techniques include image-guided intra-articular steroid injections, radiofrequency neurotomy, cryotherapy, prolotherapy, and fusion. There is a paucity of controlled trial son any treatment measure.

#### References

1. Mixer, W.J., Barr, J.S.: Rupture of the Intervertebral Disc with Involvement of the Spinal Canal. *New England Journal Med* 1934; 211:210-215.
2. Fortin, J.D., Falco, F., Washington, W.: Three Pathways Between the Sacroiliac Joint and Neural Structures Exist. *Am J Neuroradiol* 1999; 20:1429-1434.
3. Fortin JD, Tolchin R.: Sacroiliac Arthrograms and Post-Arthrography/CT. *Arch Phys Med Rehab* 1993; 74:1259.
4. Fortin, J.D.: The Sacroiliac Joint: A New Perspective. *J Back Musculoskeletal Rehabil* 1993; 3:31-43.
5. Fortin, J.D., Sehgal, N.: Sacroiliac Joint Injection and Arthrography with Imaging Correlation. In Lennard TA (ed.), *Pain Procedures in Clinical Practice*. Hanley & Beifus. Philadelphia, 2000, pp 265-273.
6. Fortin, J.D., Pier, J., Falco, F.: Sacroiliac Joint Injection: Pain Referral Mapping and Arthrographic Findings. In Vleeming A, Mooney V, Dorman T, Snijders C, Stoeckart R (eds), *Movement, Stability, and Low back Pain: The essential Role of the Pelvis*. Churchill Livingstone, New York, 1997, pp 271-285.
7. Ebraheim, N.A., Mekhail, A.O., Wiley, W.F.: Radiology of the Sacroiliac Joint. *Spine* 1997; 22: 869-876.

8. Vleeming, A., Stoeckart, R., Volkers, ACW: Relation Between Form and Function in the Sacroiliac Joint (1); The Clinical Anatomical Aspects. (II): The Biomechanical Aspects. Spine 1990; 15: 130-132.
9. Bowen, V., Cassidy, J.D.: Macroscopic and Microscopic Anatomy of the Sacroiliac Joint from Embryonic Life Until the Eight Decade. Spine 1981; 6:620-628.
10. Gunterberg, B., Romanus, B., Stener, B.: Pelvic Strength After Major Amputation of the Sacrum: An Experimental Approach. Acta Orthop Scand 1976; 47:635-642.
11. Miller, JAA, Schultz, A.M., Anderson, G.SJ: Loading Displacement Behaviors of Sacroiliac Joints. J Orthop Res 1987; 5:92-101.
12. Vleeming, A., Pool-Goudzwaard, A.L., Stoeckart, E.: The Posterior Layer of the Thoracolumbar Fascia. Spine 1995; 20:753-758.
13. Fortin, J.D., Kissling, R.O., O'Connor, B.L.: Sacroiliac Joint Innervation and Pain. Am J Orthop 1999; 12:687-690.
14. Grob, K.R., Neuhuber, W.L., Kissling, R.O.: Die Innervation des Sacroiliacalgelenkes beim Menschen. Z Rheumatol 1995; 54: 117-122.
15. Fortin, J.D., Dyer, A.P., West, S.: Sacroiliac Joint: Pain Referral Maps Upon Applying a New Injection/ Arthrography Technique – Part I: Asymptomatic Volunteers. Spine 1994; 19: 1475-1482.
16. Fortin, J.D., Vilensky, J.A., Merkel, G.J.: Can the Sacroiliac Joint Cause Sciatica? Pain Physician 2003; 6:269-271.
17. Vilensky, J.A., O'Connor, B.L., Fortin, J.D.: Histologic Analysis of Neural Elements in the Human Sacroiliac Joint. Spine 2002; 27: 1202-1207.
18. Fortin, J.D., Falco, F.: Enigmatic Causes of Spine Pain in Athletes. Phys Med Rehabil State Art Rev 1977; 11: 445-464.
19. Fortin, J.D., Aprill, C.N., Ponthieux, B.: Sacroiliac Joint: Pain Referral Maps Upon Applying a New Injection/ Arthrography Technique – Part II: Clinical Evaluation. Spine 1994; 19:1483-1489.
20. Fortin, J.D., Falco, FJE: The Fortin Finger Test: An Indicator of Sacroiliac Pain. Am J Orthop 1997; 26: 477-480.

Additional references to be posted on the website.

## KRIS C. P. VISSERS, MD, PHD, FIPP

### BIOGRAPHICAL SKETCH

Professor Vissers is an anesthesiologist, professor in Pain and Palliative Medicine and chairman of the Academic Center of Pain and Palliative Medicine of the Radboud University Nijmegen Medical Centre in the Netherlands. He is also scientific chairman of the Department of Anesthesiology, Pain and Palliative Medicine of this University.

### EPIDURAL INJECTIONS

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- The indications and rationale for epidural application of corticosteroids
- The interlaminar and transforaminal approach of the epidural space at the lumbar and cervical region
- The evidence for epidural injections (interlaminar and transforaminal) in the cervical region
- The evidence for epidural injections (interlaminar, transforaminal and causal) in the lumbar region
- The side effects and complications reported in the literature at cervical and lumbar level
- Explain measures to be taken to perform epidural injections in the safest way possible.

#### Key points

- Cervical radicular pain is defined as pain perceived as arising in the arm caused by irritation of a cervical spinal nerve or its roots
- Lumbosacral radicular pain is characterized by a radiating pain in one or more lumbar or sacral dermatomes; it may or may not be accompanied by other radicular irritation symptoms and/or symptoms of decreased function.
- The rationale for epidural corticosteroid injections is to administer the anti-inflammatory compound as close as possible to the inflamed nerve root.
- For the interlaminar administration a midline approach is used. The injection is made in the posterior compartment. The spread towards the ventral part of the epidural space could be limited
- The transforaminal administration of corticosteroids aims at injecting the corticosteroid directly onto and around the nerve root.
- There are no comparative trials investigating interlaminar and transforaminal injections at cervical level.
- An older randomized trial compares interlaminar injection with intramuscular injections and finds the former to yield superior results. One more recent study investigates the added effect of morphine and another trial compares a single injection with continuous infusion.
- A randomized controlled trial compares the effect of cervical transforaminal injection of corticosteroid plus local anesthetic with local anesthetic alone or saline and found no difference in outcome between the three groups.
- The effect of lumbar epidural interlaminar corticosteroid injections was studied in systematic reviews. These conclude that this technique provides pain relief for a relative short duration.
- Lumbar transforaminal epidural corticosteroid administration was studied in randomized controlled trials. One trial demonstrated that this treatment is effective for the management of contained herniations.
- Cervical transforaminal epidural corticosteroid administration has resulted in several cases of serious unexpected and unexplained complications and therefore is not recommended.
- Anatomic variations of the vascular supply in the cervical area supports the statement that there is no safe area for needle placement in the cervical foramina. The side effects and complications reported with the use of the interlaminar cervical epidural corticosteroid injections are in the vast

majority of the cases minor and transient. There are however two reports of permanent damage to the spinal cord which occurred in sedated patients.

- In the lumbar region side effects of interlaminar corticosteroid administration are comparable to those reported for the cervical area. With transforaminal epidural administration cases of neurological complications have been reported.
- To reduce the risk of complications epidural injections should always be performed under radiographic control with use of contrast medium. When optimal needle placement is achieved a test dose of local anesthetic and an observation period of 60 sec is recommended.
- The use of non-particulate corticosteroid is preferred by some.

#### Recommended literature

\* of special interest

\*\* of outstanding interest

1. \*\* Abbasi A, Malhotra G, Malanga G, Elovic EP, Kahn S. Complications of interlaminar cervical epidural steroid injections: a review of the literature. *Spine* 2007;32:2144-2151.
2. Abdi S, Datta S, Trescot AM, Schultz DM, Adlaka R, Atluri SL, Smith HS, Manchikanti L. Epidural steroids in the management of chronic spinal pain: a systematic review. *Pain Physician* 2007;10:185-212.
3. \* Ackerman WE, 3rd, Ahmad M. The efficacy of lumbar epidural steroid injections in patients with lumbar disc herniations. *Anesth Analg* 2007;104:1217-1222.
4. \* Anderberg L, Annertz M, Persson L, Brandt L, Saveland H. Transforaminal steroid injections for the treatment of cervical radiculopathy: a prospective and randomised study. *Eur Spine J* 2007;16:321-328.
5. \* Karppinen J, Malmivaara A, Kurunlahti M, Kyllonen E, Pienimäki T, Nieminen P, Ohinmaa A, Tervonen O, Vanharanta H. Periradicular infiltration for sciatica: a randomized controlled trial. *Spine* 2001a;26:1059-1067.
6. \* Karppinen J, Ohinmaa A, Malmivaara A, Kurunlahti M, Kyllonen E, Pienimäki T, Nieminen P, Tervonen O, Vanharanta H. Cost effectiveness of periradicular infiltration for sciatica: subgroup analysis of a randomized controlled trial. *Spine* 2001b;26:2587-2595.
7. \* Koes B, Scholten RJ, Mens JMA, Bouter LM. Efficacy of epidural steroid injections for low-back pain and sciatica: a systematic review of randomized clinical trials. *Pain* 1995;63:279-288.
8. \*\* Malhotra G, Abbasi A, Rhee M. Complications of transforaminal cervical epidural steroid injections. *Spine* 2009;34:731-739.
9. \* McQuay H, Moore A. An evidence-based resource for pain relief. Oxford University Press 1998.
10. Peloso P, Gross A, Haines T, Trinh K, Goldsmith CH, Burnie S. Medicinal and injection therapies for mechanical neck disorders. *Cochrane Database Syst Rev* 2007(3):CD000319.
11. Riew KD, Park JB, Cho YS, Gilula L, Patel A, Lenke LG, Bridwell KH. Nerve root blocks in the treatment of lumbar radicular pain. A minimum five-year follow-up. *J Bone Joint Surg Am* 2006;88:1722-1725.
12. Riew KD, Yin Y, Gilula L, Bridwell KH, Lenke LG, Laurusen C, Goette K. The effect of nerve-root injections on the need for operative treatment of lumbar radicular pain. A prospective, randomized, controlled, double-blind study. *J Bone Joint Surg Am* 2000;82-A:1589-1593.
13. Stav A, Ovadia L, Sternberg A, Kaadan M, Weksler N. Cervical epidural steroid injection for cervicobrachialgia. *Acta Anaesthesiol Scand* 1993;37:562-566.
14. \*\* Van Boxem K, Cheng J, Patijn J, van Kleef M, Lataster A, Mekhail N, Van Zundert J. 11. Lumbosacral radicular pain. *Pain Pract* 2010;10:339-358.
15. \*\* Van Zundert J, Huntoon M, Patijn J, Lataster A, Mekhail N, van Kleef M. 4. Cervical radicular pain. *Pain Pract* 2009;10:1-17.

## JOSEPH D. FORTIN, MD

### IMAGING FOR INTERVENTIONAL PAIN MANAGEMENT 2011

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- Selecting and applying appropriate imaging studies in a clinical context.
- Utilizing a structure and function approach to interpreting imaging studies.
- Limitations of various imaging studies, as well as the combined benefits of specific modalities.

#### Key Points

- Imaging studies represent a key diagnostic tool for the pain management physician; yet lack of a concerted exposure to integrate these modalities in pain management training programs is widely recognized.
- Understanding the indications for the various imaging modalities and how alterations in imaging anatomy may reflect an injury mechanism provides the pain management physician a powerful compliment to their diagnostic armamentarium.
- Plain films still fulfill the basic roles within our diagnostic scope for screening trauma victimcases; when ruling out instability, examining for benign bone conditions and providing a comparative or complimentary study to magnetic resonance imaging, computerized tomography, or ultrasound.
- CT has higher spatial resolution than MRI and consequently provides superior contour resolution for depicting contour changes and cortical bone margins.
- For superficial structures ultrasound has resolution that exceeds MRI and consequently is especially effective in examining most parts of the peripheral musculoskeletal system. It is also arguably the most dynamic and portable imaging modality. Ultrasound is limited by being operator dependent, having a small field of view, suboptimal tissue contrast, and requiring low frequency at the cost of resolution for deep tissue penetration.
- Scintigraphy is very sensitive for AVN, occult fractures, and most osseous tumors and infections. Unfortunately, it is radiologically non-specific.
- MRI is the study of choice for evaluating soft tissue pathology including the intrathecal space and it's neurovascular contents. It is also an elegant modality for staging degenerative, inflammatory, or traumatic marrow space changes.

#### References

1. Fortin, J.D.: An Algorithm for Understanding Spine Imaging. *Pain Physician*. 2002, 5:102-9.
2. Fortin, J.D., Weber, E.C.: The Fundamentals of Musculoskeletal Imaging. *Pain Physician*. 2004 7: 149-160,
3. Fortin, J.D., Wheeler, M.T.: Imaging in Lumbar Spinal Stenosis. *Pain Physician*. 2004 7: 133-139.
4. Fortin, J.D., Weber, E.C.: Imaging of Whiplash Injuries. *Spine: State of the Art Reviews* 1998 12:419-436,
5. Kane, D., Balint, P.V., Sturrock, R., Grassi, W.: Musculoskeletal Ultrasound – A State of the Art Review in *Rheumatology: Part 1 Current Controversies and Issues in the Development of Musculoskeletal Ultrasound in Rheumatology*. 2004 43: 823-828
6. Grassi, W., Filippucci, E., Busilacchi, P.: Musculoskeletal Ultrasound: Best Practice and Research Clinical *Rheumatology* 2004 Vol. 18, No 6, 813-824

## RAY M. BAKER, MD, FIPP

### BIOGRAPHICAL SKETCH

Dr. Baker is a Clinical Professor of Anesthesiology (adjunct) at the University of Washington. He is the Immediate Past President of the North American Spine Society and the incoming President of the International Spine Intervention Society.

### CONTROVERSIES IN THE DIAGNOSIS OF PAINFUL LUMBAR DISC DEGENERATION 2011

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- The definition and key features of a diagnostic criterion standard
- The role of advanced imaging in the diagnosis of painful lumbar disc degeneration (DD)
- The role of the history and physical examination in the diagnosis of painful lumbar DD
- The role of provocation discography in the diagnosis of painful lumbar DD
- Controversies in the diagnosis of painful lumbar DD, including:
  - The limitations of provocation discography
  - The potential risks of provocation discography
  - The link between provocation discography and expensive treatments, especially lumbar fusion
- The potential role of novel technologies in the diagnosis of painful lumbar DD

#### Key Points

- Utilizing the Sackett and Haynes definition of a criterion standard, there exists no Gold Standard for the diagnosis of painful lumbar DD.
- Advanced imaging is helpful in excluding patients with normal MRIs from the diagnosis of painful lumbar DD.
- Compared with provocation discography, MRI loss of signal intensity, high intensity zones, and end-plate (Modic) changes are modestly helpful at best (+LR 3) in diagnosing patients with painful lumbar DD.
- The history and physical examination are of limited utility in diagnosing patients with painful lumbar DD.
- Provocation discography is still a controversial test for painful lumbar DD. There is no ability to compare it with a criterion standard, there is a high false positive rate in patients with certain co-existing conditions, and it has been linked to poor outcomes from lumbar fusion surgery. Recent studies have also raised the possibility that disc puncture might accelerate disc degeneration.
- Limited evidence supports novel technologies in the diagnosis of painful lumbar DD, including MR Spectroscopy.

#### References

1. Carragee EJ, Chen Y, Tanner CM, et al. Can discography cause long-term back symptoms in previously asymptomatic subjects? *Spine* 2000;25:1803–8.
2. Carragee EJ, Chen Y, Tanner CM, et al. Provocative discography in patients after limited lumbar discectomy: a controlled, randomized study of pain response in symptomatic and asymptomatic subjects. *Spine* 2000; 25:3065–71.
3. Carragee EJ, Tanner CM, Khurana S, et al. The rates of false-positive lumbar discography in select patients without low back symptoms. *Spine* 2000;25: 1373–80; discussion 1381.
4. Carragee E, Barcohana B, Alamin TF, et al. Prospective controlled study of the development of lower back pain in previously asymptomatic subjects undergoing experimental discography. *Spine* 2004;29:1112–7.
5. Carragee EJ, Alamin TF, Miller JL, et al. Discographic, MRI and psychosocial determinants of low back pain disability and remission: a prospective study in subjects with benign persistent back pain. *Spine J* 2005;5:24–35.

6. Carragee EJ, Don AS, Hurwitz EL, et al. Does discography cause accelerated progression of degeneration changes in the lumbar disc: a ten-year matched cohort study. *Spine* 2009;34:2338–45.
7. Hancock MJ, Maher CG, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J* (2007) 16:1539–1550
8. Hägg O, Fritzell P, Ekselius L, et al. Predictors of outcome in fusion surgery for chronic low back pain. A report from the Swedish Lumbar Spine Study. *Eur Spine J*. 2003 Feb;12:22-33.
9. Jarvik JG, Hollingworth W, Heagerty PJ, et al. Three-Year Incidence of Low Back Pain in an Initially Asymptomatic Cohort. *Spine* 2005. 30:1541–1548.
10. Keshari KR, Lotz JC, Kurhanewicz J, and Majumdar S. Correlation of HR-MAS Spectroscopy Derived Metabolite Concentrations With Collagen and Proteoglycan Levels and Thompson Grade in the Degenerative Disc. *Spine* 2005;30:2683–2688.
11. Korecki CL, Costi JJ, Iatridis JC. Needle puncture injury affects intervertebral disc mechanics and biology in an organ culture model. *Spine* 2008;33:235–41.
12. Nassr A, Lee JY, Bashir RS, et al. Does incorrect level needle localization during anterior cervical discectomy and fusion lead to accelerated disc degeneration? *Spine* 2009;34:189–92.
13. Sackett DL, Haynes RB. The architecture of diagnostic research. *BMJ* 2002;324:539–41.

## BIOGRAPHICAL SKETCH

Dr. Bruni is in charge of the Clinical Trials Unit, Department of Biomedicine - Division of Rheumatology, University of Florence, Italy

## JOINT PAIN AND ITS TREATMENT WITH BIOLOGICAL AGENTS AND DRUGS

### Objectives

Upon completion of this presentation attendees will be able to discuss:

- The importance of pain in rheumatic diseases
- The role of cytokines in inflammation and pain
- How to assess joint pain in daily practice
- How to assess disease activity in Rheumatoid Arthritis
- Biological therapy and Target therapy
- Effects of TNF- $\alpha$  inhibitors on pain during RCTs
- The comparison between Biological drugs and DMARDs
- Goals of therapy in Rheumatic diseases

### Key Points

- Pain is one of the main features of rheumatic diseases and its management is an area of increasing research.
- Pain in rheumatic diseases is strictly connected with inflammation, whose pathogenesis depends on many cytokines as TNF- $\alpha$ , IL-1 $\beta$  and IL-6, which have also a very important role in the maintenance of pain and in the CNS.
- Old drugs like DMARDs are able to control mainly inflammation, with a minor effect on disability and bone damage.
- It's important to assess Pain in daily practice and it is also a parameter of the Disease Activity Score (DAS), which is the best index to assess disability too.
- Biological drugs are produced using biotechnology and are directed against specific cytokines or molecular pathways: this is the so-called Target Therapy.
- The first biological drugs were TNF- $\alpha$  inhibitors, which proved to be effective in reducing pain, improving quality of life and managing disease activity in RA and other rheumatic and non-rheumatic diseases, as shown by many RCTs.
- In rheumatological diseases the sooner the therapy is started, the better the disease activity is controlled, as patients seem to be more prone to favourable treatment outcome during the very start of the disease.

### References

1. Schaible HG et al. The role of proinflammatory cytokines in the generation and maintenance of joint pain. *Ann. N.Y. Acad. Sci.* 2010; 1193:60–69.
2. Boettger MK et al. Antinociceptive effects of tumor necrosis factor alpha neutralization in a rat model of antigen-induced arthritis: evidence of a neuronal target. *Arthritis Rheum* 2008;58:2368–78.
3. Kirwan JR. Links between radiological change, disability, and pathology in rheumatoid arthritis. *J Rheumatol.* 2001;28:881-886.
4. Van Gestel AM, Prevoo MLL, van't Hof MA, et al. Development and validation of the European League Against Rheumatism response criteria for rheumatoid arthritis. *Arthritis Rheum* 1996; 39:34-40.
5. Van der Heijde et al. Validity of single variables and composite indices for measuring disease activity in *Ann Rheum Dis* 1992;51:177-81.
6. Maini RN et al. Sustained improvement over two years in physical function, structural damage, and signs and symptoms among patients with rheumatoid arthritis treated with infliximab and methotrexate. *Arthritis Rheum* 2004; 50: 1051-65.
7. Weinblatt ME et al. Adalimumab, a fully human anti-tumor necrosis factor alpha monoclonal antibody, for the treatment of rheumatoid arthritis in patients taking concomitant methotrexate: the ARMADA trial. *Arthritis Rheum.* 2003;48:35-45.

8. Kekow J et al. Patient-reported outcomes improve with etanercept plus methotrexate in active early rheumatoid arthritis and the improvement is strongly associated with remission: the COMET trial. *Ann Rheum Dis* 2010 Jan; 69: 222-5.
9. Russell AS et al. Abatacept improves both the physical and mental health of patients with rheumatoid arthritis who have inadequate response to methotrexate treatment. *Ann Rheum Dis* 2007;66: 189-94.
10. Keystone E et al. Improvement in patient-reported outcomes in a rituximab trial in patients with severe rheumatoid arthritis refractory to anti-tumor necrosis factor therapy. *Arth Rheum* 2008;59:785-93.
11. Furst D, Window of opportunity, *J Rheumatol* 2004;31:1677-9.

## LUMBOSACRAL SPINAL ENDOSCOPY ROUND TABLE

**SCIENCE BACKGROUND: JAMES E. HEAVNER, DVM, PHD, FIPP (HON)**

**CLINICAL PERSPECTIVE: HEMMO A. BOSSCHER, MD, FIPP**

**APPLICATION TO ARACHNOIDITIS: JAN PETER WARNKE, MD**

### Objectives

Upon completion of this round table presentation attendees will be able to discuss

- Key developments leading to the use of flexible endoscopes in the spinal canal
- Primary reasons for performing spinal endoscopy
- Indications and techniques for performing epiduroscopy and thecaloscopy
- How patients benefit from spinal canal endoscopy

### Key Points

- Advent of small, flexible fiberscopes and suitable light sources, as well as techniques to introduce scopes into the spinal canal is the foundation for spinal endoscopy
- Spinal canal endoscopy provides direct viewing of structures within the canal thereby aiding in determination of abnormalities involved in chronic low back pain and/or pain radiating to the lower extremities
- Spinal canal endoscopy allows direct therapeutic intervention while viewing sources of pain
- Spinal canal endoscopy may identify causes of pain that cannot be determined by physical examination and imaging (CT scan, MRI)
- Major surgical intervention can often be avoided by using minimally invasive spinal canal endoscopy.

### References

1. Heavner JE, Bosscher HA, Anderson S. Epiduroscopy. In: *Interventional Pain Management: Image Guided Procedures*, 2nd ed. P Raj, et al, eds. Philadelphia: Saunders Elsevier, 2008: 529-536.
2. Heavner JE, Bosscher HA, Wachtel M. Cell types obtained from the epidural space of patients with low back pain/radiculopathy. *Pain Practice*. 2009; 9: 167-172.
3. Bosscher HA, Heavner JE. Incidence and severity of epidural fibrosis after back surgery: An endoscopic study. *Pain Practice*. 2010; 10: 18-24.
4. Mourgela A, Sakellaropoulos A, Anagnostopoulou A, Warnke JP. The dimensions of the sacral spinal canal in thecaloscopy: A morphometric MRI study. *Neuroanatomy*. 2009; 8: 1-3.
5. Di Ieva A, Barolat G, Tschabitscher M, Rognone E, Aimar E, Gaetani P, Tancioni F, Lorenzetti M, Crotti FM, Rodriguez Baena R, Warnke JP. Lumbar arachnoiditis and thecaloscopy: Brief review and proposed treatment algorithm. *Cen Eur Neurosurg*. 2010; 71: 207-212.

## COLD ALLODYNIA, NERVE ENTRAPMENT, RADICULOPATHIES AND ILLUSIONS: THE SCIENCE AND THE CLINICAL

**THE SCIENCE: JAMES E. HEAVNER, DVM, PHD, FIPP (HON)**

**THE CLINICAL: GABOR B. RACZ, MD, FIPP**

### Objectives

Upon completion of this round table presentation attendees will be able to discuss

- Cold allodynia and hyperalgesia as frequent clinical findings in patients with neuropathic pain
- Mechanisms and nerve pathways implicated in cold allodynia and hyperalgesia
- Case examples of cold allodynia and hyperalgesia in patients with neuropathic pain
- Therapies – what works, what does not work and why

### Key Points

- Cold allodynia and hyperalgesia are associated with injuries to both the central and peripheral nervous systems
- In general, little is known about mechanisms of cold allodynia and hyperalgesia. NMDA-receptor mediated central sensitization is involved in cold hyperalgesia but other mechanisms are also present
- The normal human brain can be tricked to perceive cold allodynia (thermal grill illusion) which might provide clues to how nerve injury produces cold allodynia and hyperalgesia
- Sympathetic block may relieve cold allodynia/hyperalgesia, but may not be successful if the block is not complete

### References

1. Jorum E, Warncke T, Stubhaug A. Cold allodynia and hyperalgesia in neuropathic pain: The effect of N-methyl-D-aspartate (NMDA) receptor antagonist ketamine – a double-blind, cross-over comparison with alfentanil and placebo. *Pain*. 2003; 101: 229-235.
2. Costigan M, Scholz J, Woolf CJ. Neuropathic pain: A maladaptive response of the nervous system to damage. *Annu Rev Neurosci*. 2009; 32: 1-32.
3. Heavner JE, Calvillo O, Racz GB. Thermal grill illusion and complex regional pain syndrome I (reflex sympathetic dystrophy). *Reg Anesth*. 1997; 22: 257-9.
4. Heavner JE, Willis WD. Pain pathways: Anatomy and physiology. In: *Practical Management of Pain*, 3rd ed. P Raj, ed. St. Louis: Mosby, 20

## ANDREA TRECOT, MD, FIPP

### BIOGRAPHICAL SKETCH

Dr. Andrea Trecot is a private practice physician in St. Augustine, Florida.

### PATHOPHYSIOLOGY AND PHARMACOLOGIC MANAGEMENT OF NEUROPATHIC PAIN

#### Objectives

- Upon completion of this presentation, attendees will be able to discuss:
- Some of the common causes of peripheral neuropathy
- Some of the common pharmacologic and nonpharmacologic treatments of peripheral neuropathy
- Some of the new directions in the treatment of peripheral neuropathy

#### Key points

- Peripheral neuropathy is not a homogeneous process, but rather encompasses a large, diverse group of nerve pathologies.
- There are multiple causes of peripheral neuropathy.
- Although diabetes is the most common cause of peripheral neuropathy, most cases of peripheral neuropathy never have a defined etiology.
- Most of the treatments are pharmacologic, but there are increasingly options for interventional treatment

## KRIS C. P. VISSERS, MD, PHD, FIPP

### PHARMACOLOGICAL MANAGEMENT OF CANCER PAIN

#### Objectives

At the end of this presentation attendees will be able to discuss:

- The value of following the WHO pain ladder in the management of cancer pain
- The role of adjuvant treatment such as antidepressants and anti-epileptics
- The appropriate use of breakthrough medication
- The control of side effects and complications
- The role of palliative cancer treatments such as chemotherapy and bisphosphonates
- The right dose calculation for an opioid switch

#### Key points

- The management of cancer pain requires a multidisciplinary and multifactorial approach. The treatment may consist of tumor oriented interventions supplemented with symptomatic pain management.
- The most important step in reaching an adequate pain control in patients with cancer is a good understanding of the underlying pain mechanism, in order that the right treatment can be chosen
- The first step in pain treatment is antitumor therapy by the medical oncologist if indicated
- In case of pain caused by bone metastases, bisphosphonates may be considered.
- Basic pharmacological treatment aims at achieving constant steady state plasma levels, which is preferentially achieved by long-acting formulations.
- Patients should be instructed to use immediate release preparations for control of breakthrough pain.
- The use of step two of the WHO pain ladder is no longer recommended.
- Only weak recommendation can be used to support combination step three opioid therapy.
- There is evidence to support the efficacy and tolerability of hydromorphone for moderate to severe cancer pain as an alternative to morphine and oxycodone.
- Alternative administration routes to oral opioids such as parenteral, transdermal and rectal administration may be considered. There is no significant difference in efficacy or side effects between administration routes.
- There is a weak recommendation for using spinal opioids in adult patients with cancer.
- In case of tolerance or severe side effects on opioid analgesics, opioid switch is recommended.
- When opioid therapy is started it is recommended to prescribe osmotic laxatives.
- Vomiting and nausea may be controlled by prokinetics and dopamine antagonists
- The direct clinical evidence in cancer-related pain and renal impairment is insufficient to allow formulation of guidelines. The risk of opioid use is stratified according to the activity of opioid metabolites and potential for accumulation. Fentanyl, alfentanil and methadone are the least likely to cause harm when used appropriately. Morphine may be associated with toxicity in patients with renal impairment.
- The role of adjuvant therapy (e.g. antidepressants and anti-epileptics) has become evident when neuropathic characteristics are present.
- Side effects of the central nervous system caused by opioid analgesics, such as sedation and hallucinations, may wear off over time.
- Delirium, myoclonus and hyperalgesia may be controlled by opioid switching
- For the management of opioid intoxication opioid antagonists such as naloxone are recommended.
- Recently, opioid antagonists, that do not cross the blood brain barrier have been developed. Simultaneous use of those antagonists and opioids may prevent gastro-intestinal side effects.

## Recommended literature

\* of special interest

\*\* of outstanding interest

1. \*\* Portenoy, R. K. (2011). Treatment of cancer pain. *Lancet* 377, 2236-2247.

## Chemotherapy

2. Anderson H, Hopwood P, Stephens RJ, Thatcher N, Cottier B, Nicholson M, Milroy R, Maughan TS, Falk SJ, Bond MG, Burt PA, Connolly CK, McIlmurray MB, Carmichael J. Gemcitabine plus best supportive care (BSC) vs BSC in inoperable non-small cell lung cancer—a randomized trial with quality of life as the primary outcome. UK NSCLC Gemcitabine Group. *Non-Small Cell Lung Cancer. Br J Cancer* 2000;83:447-453.
3. Dancey J, Shepherd FA, Gralla RJ, Kim YS. Quality of life assessment of second-line docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy: results of a prospective, randomized phase III trial. *Lung Cancer* 2004;43:183-194.
4. Osoba D, Tannock IF, Ernst DS, Neville AJ. Health-related quality of life in men with metastatic prostate cancer treated with prednisone alone or mitoxantrone and prednisone. *J Clin Oncol* 1999;17:1654-1663

## Biphosphonates

5. Djulbegovic B, Wheatley K, Ross J, Clark O, Bos G, Goldschmidt H, Cremer F, Alsina M, Glasmacher A. Bisphosphonates in multiple myeloma. *Cochrane Database Syst Rev* 2002(3):CD003188.
6. Pavlakis N, Schmidt R, Stockler M. Bisphosphonates for breast cancer. *Cochrane Database Syst Rev* 2005(3):CD003474.
7. Wong R, Wiffen PJ. Bisphosphonates for the relief of pain secondary to bone metastases. *Cochrane Database Syst Rev* 2002(2):CD002068.
8. Yuen KK, Shelley M, Sze WM, Wilt T, Mason MD. Bisphosphonates for advanced prostate cancer. *Cochrane Database Syst Rev* 2006(4):CD006250.

## Weak opioids

9. Marinangeli F, Ciccozzi A, Leonardis M, Aloisio L, Mazzei A, Paladini A, Porzio G, Marchetti P, Varrassi G. Use of strong opioids in advanced cancer pain: a randomized trial. *J Pain Symptom Manage* 2004;27:409-416.
10. \* Maltoni M, Scarpi E, Modonesi C, Passardi A, Calpona S, Turriziani A, Speranza R, Tassinari D, Magnani P, Saccani D, Montanari L, Roudnas B, Amadori D, Fabbri L, Nanni O, Raulli P, Poggi B, Fochessati F, Giannunzio D, Barbagallo ML, Minnotti V, Betti M, Giordani S, Piazza E, Scapaticci R, Ferrario S. A validation study of the WHO analgesic ladder: a two-step vs three-step strategy. *Support Care Cancer* 2005;13:888-894.

## Strong opioids

11. \* Nicholson AB. Methadone for cancer pain. *Cochrane Database Syst Rev* 2004(2):CD003971.
12. \* Fallon, M. T., and Laird, B. J. (2011). A systematic review of combination step III opioid therapy in cancer pain: An EPCRC opioid guideline project. *Palliat Med* 25, 597-603.
13. Radbruch, L., Trottenberg, P., Elsner, F., Kaasa, S., and Caraceni, A. (2011). Systematic review of the role of alternative application routes for opioid treatment for moderate to severe cancer pain: An EPCRC opioid guidelines project. *Palliat Med* 25, 578-596.
14. Pigni, A., Brunelli, C., and Caraceni, A. (2011). The role of hydromorphone in cancer pain treatment: a systematic review. *Palliat Med* 25, 471-477.

## Opioid rotation

15. \*\* Vissers KC, Besse K, Hans G, Devulder J, Morlion B. Opioid Rotation in the Management of Chronic Pain: Where Is the Evidence? *Pain Pract* 2010.

## Breakthrough medication

16. Hanks GW, Conno F, Cherny N, Hanna M, Kalso E, McQuay HJ, Mercadante S, Meynadier J, Poulain P, Ripamonti C, Radbruch L, Casas JR, Sawe J, Twycross RG, Ventafridda V. Morphine and alternative opioids in cancer pain: the EAPC recommendations. *Br J Cancer* 2001;84(5):587-593.

## Adjuvant analgesics

17. Lussier D, Huskey AG, Portenoy RK. Adjuvant analgesics in cancer pain management. *Oncologist* 2004;9:571-591.

## Opioid side effects

18. McNicol E, Horowicz-Mehler N, Fisk RA, Bennett K, Gialeli-Goudas M, Chew PW, Lau J, Carr D. Management of opioid side effects in cancer-related and chronic noncancer pain: a systematic review. *J Pain* 2003;4:231-256.
19. King, S., Forbes, K., Hanks, G., Ferro, C., and Chambers, E. (2011). A systematic review of the use of opioid medication for those with moderate to severe cancer pain and renal impairment: A European Palliative Care Research Collaborative opioid guidelines project. *Palliat Med* 25, 525-552.

## LORAND EROSS, MD, PHD, FIPP

### BIOGRAPHICAL SKETCH

Dr. Lorand Eross is the director of Functional Neurosurgical Program and head of the Functional Neurosurgical Department at the National Institute of Neuroscience in Budapest. He is a board certified neurologist and neurosurgeon. He has got his PhD degree at Semmelweis University, Faculty of Medicine in 2010. His main interest is epilepsy surgery, movement disorder surgery, pain treatment, spasticity, intraoperative neuromonitoring and neuromodulation. He is teaching at Semmelweis University at the Faculty of Medicine and at Pazmany Peter University Faculty of Information Technology. His research activity is in vitro and in vivo electrophysiological investigational methods in epilepsy.

### CURRENT NEUROSURGICAL APPROACHES TO THE TREATMENT OF PAIN

#### Learning objectives:

This summary focuses exclusively on neurosurgical procedures against pain. SCS and peripheral nerve stimulation will be discussed by other authors.

#### Key Points:

The neurosurgical treatment of pain is divided into two subgroups: ablative and neuroaugmentative therapies.

**Ablative procedures** include all types of surgical interventions, when an irreversible action is taken to stop pain.

**Neurolysis:** separation of a peripheral nerve from the surrounding structures to which is adherent. The use of internal neurolysis is clearly necessary in dissecting an injured nerve for interfascicular nerve graft or to evaluate a neuroma-in-continuity.

**Trigeminal neurectomy:** Peripheral trigeminal neurectomy can be useful in elderly debilitated patients who cannot undergo more substantive procedure for V1 division neuralgia. 50-60% of trigeminal neuropathic pain cases are successfully treated with neurectomy.

**Dorsal Root ganglionectomy and Dorsal Rhizotomy (DR):** The largest series of DR in cancer pain was published in 1982 by Sindou and Lapras, success rate was 47% in a series of 585 patients.

**Sympathectomy:** Currently surgical sympathectomy is reserved for treating hyperhydrosis, sympathetically maintained pain and limited cases of vasculitis (i.e. Raynaud's syndrome). The success rate of sympathectomy in the literature after 1990 ranges from 65% to 100%.

**Dorsal Root Entry Zone lesioning:** Indications for drezotomy includes 1.Cancer pain that is limited in extent (e.g.: Pancoast syndrome), 2.Persistent neuropathic pain, 3.Disabling hyperspasticity, especially when associated with pain. Surgery in the DREZ must be considered within the frame of all the methods belonging to the armamentarium of pain surgery.

**Midline myelotomy:** Gildenberg and Hirschberg (1984) performed myelotomy for visceral pain with excellent results in 8 out of 12 patients. Punctuate midline myelotomy after laminectomy at T8 level for malignant visceral pain found efficient by Nauta et al. (2000). This technique has limited indication today.

**Anterior Cordotomy:** The ideal candidates for Percutan Cordotomy (PC) are cancer patients with unilateral localized pain if the primary malignant disease is under control. The initial success rate of 3742 cases collected by Lorenz was 75 to 96%.

**Percutaneous extralemniscal myelotomy:** Indicated in cancer patients with pelvic or lower trunk or lower extremity pain. Kanpolat reported 15 cases, with rectal, pancreatic, colon, renal tumors without complication rate. 6 of the 15 patient had complete 5 of 15 cases had partial pain relief.

**Mesencephalotomy:** Amano in 1998 reported 76% long-term pain relief in patients with central and deafferentation pain with an overall morbidity of 4%. No recent report of this procedure in practice.

**Medial Thalamotomy (MT):** MT is capable of alleviating neuropathic and nociceptive pain and has the advantage of low morbidity. Medial thalamotomy in any nucleus is more effective in relieving nociceptive than neuropathic pain and those results are modest: 46% relief of nociceptive usually cancer



**Unmatched Physician & Patient Support**

- Cadaver Workshops
- Procedure Reference Literature
- Sponsor of Clinical Research
- Reimbursement and Physician Services

ENVISION  
DEVELOP  
EVOLVE

- Catheters for Acute, Chronic & Regional Pain
- Standard & Specialty Needles
- R-F™ Products
- Radiation Safety

For more information please call 800.866.3342  
or visit us at [epimedpain.com](http://epimedpain.com)

pain and 29% in neuropathic pain.

**Stereotactic cingulotomy:** 394 patients were reported until today, in patients with benign origin 53% was useful and 47% of non-useful. In malignant pain the result was just similar. The initial good response to cingulotomy progressively fades over time.

**Hypophysectomy:** There are few clinical report on hypophysectomy for pain in the literature since 1984. Recently some center reported on few patients gamma knife hypophysectomies with limited results.

**Percutan Radiofrequency Trigeminal Gangliolysis or Rhizotomy:** In summary of several series of RF trigeminal rhizolysis 99% of patients became pain free immediately after the procedure. In a review of 1200 patients followed 1-20 years (mean 9 years), 93 % reported excellent or good results, and 4% reported fair results because undesirable side effects, 1% reported poor results because of severe denervation dysesthesia. RF trigeminal rhizolysis is effective in primary trigeminal neuralgia. RF lesioning can effectively treat paroxysmal facial pain associated with tumors and multiple sclerosis.

**Percutan Retrogasserian Glycerol Rhizotomy (PRGR)** PRGR is a useful minimal invasive technique in trigeminal neuralgia when MVD is not possible. Long term pain control (7 years) was 85%; the 11 years follow up in Lundsford series showed 77% pain relief.

**Microvascular decompression (MVD)** for Trigeminal Neuralgia: Jannetta reported a total success rate of 88% at 1 year and 74% at 10 year follow up. MVD is the treatment of choice for patient with typical trigeminal neuralgia, with MRI diagnosed neurovascular compression if the patient medical condition allow the risk of craniotomy.

**Posterior Fossa Trigeminal Rhizotomy (PFTR):** Several contemporary neurosurgeons indicate PFTR when MVD surgery or other procedures failed. In 3% of patients operated with MVD no vascular compression is found. In these cases an optional treatment strategy could be partial sectioning the nerve.

**Gamma Knife Radiosurgery for Trigeminal Neuralgia:** With this method by the end of 2010 more than 17 000 patients were treated worldwide. Approximately 75% of patients achieve good (pain free on medication) or excellent results (pain free w/o medication) within 1-8 weeks of the initial treatment.

**Neuromodulative therapy** includes only reversible neurostimulation type procedures: **Primary Motor Cortex Stimulation (MCS):** Chronic epidural MCS can control central deafferentation pain in 45-75% of cases. The best results were observed in central post-stroke pain and trigeminal neuropathy (>90%). The results improved during the last 10 years due to better targeting of the motor cortex (fMRI, neuronavigation, SSEP, intraoperative stimulation).

**Deep Brain Stimulation (DBS):** In general patients with refractory neuropathic pain should undergo paraesthesia producing stimulation, whereas those with nociceptive pain should undergo periventricular gray/periaqueductal gray matter stimulation, long-term success rate varies between 26% to 72%. The best results of DBS are in cancer pain, FBSS, cervical and brachial avulsions and peripheral neuropathy.

**Gasserian Ganglion Stimulation:** Stimulation of the gasserian ganglion presents a surgical option with atypical trigeminal pain. In a large clinical series of 182 patients 92 had more than 50% pain relief and 82 were implanted. At long-term follow up 70% of patients had 75 -100% pain relief. The most benefiter were patients with neuropathic pain after intervention of the maxillary sinus, posttraumatic facial pain, and those with severe dysesthesia after trigeminal destructive procedures.

**Key References:**

1. North RB Levy RM Consensus conference on the management of pain. Neurosurgery 1994;34
2. Burchiel K.J. Surgical management of Pain ed. Thieme Verlag 2002.
3. Lefaucheur JP, Keravel Y, Nguyen JP. Treatment of poststroke pain by epidural motor cortex stimulation with a new octopolar lead Neurosurgery, 2011 Mar; 68(1 Suppl Operative):180-7; discussion 187
4. Racz GB, Rui-Lopez R: Radiofrequency procedures Pain Pract 2006 Mar; 6(1): 46-50 Review

## LIONG LIEM, MD, FIPP

### BIOGRAPHICAL SKETCH

Dr. Liong Liem, MD, FIPP, is director of the multidisciplinary Pain Clinic in St. Antonius Hospital, Department of Anesthesiology, Intensive Care and Pain Medicine, Nieuwegein, The Netherlands.

### FACIAL PAIN AND CERVICOGENIC HEADACHE

#### Objectives

Upon completion of this presentation attendees will be able to discuss:

- The pathophysiology of and risk factors for facial pain
- The diagnostic criteria for facial pain and cervicogenic headache.
- Questions to determine the differential diagnosis of facial pain.
- Physical examination to find evidence for cervicogenic headache
- Treatment options for facial pain and cervicogenic headache if conservative therapy fails.

#### Key Points

- Persistent idiopathic facial pain (PIFP) is described as a persistent facial pain that does not have the classical characteristics of cranial neuralgias and for which there is no obvious cause. The pathophysiology is unknown.
- Forming a diagnosis is a process of elimination of other causes of facial pain and is possible if the pain is present daily and throughout all or most of the day.
- Treatment of PIFP requires a multidisciplinary approach of psychological counseling and pharmacological therapy, Amitriptyline is the primary choice. Invasive procedures can be considered if pharmacological treatment fails.
- Cervicogenic headache is characterized by unilateral headache symptoms which arise from the neck and radiates to fronto-temporal region and possible to the supraorbital region.
- Physical examination to diagnose cervicogenic headache encompasses movement tests of the cervical facet joints and soft tissues of the neck.
- Injection of the n. occipitalis major or RF treatment can be preformed if conservative treatments falls short.

#### References

1. Cornelissen P, van Kleef M, Mekhail N, Day M, van Zundert J. Persistent Idiopathic Facial Pain. Pain Practice. 9: 443-448, 2009.
2. van Suijlekom H, van Zundert J, Narouze S, van Kleef M, Mekhail N. Cervicogenic Headache. Pain Practice. 10: 124-130, 2010.
3. Bogduk N, Govind J. Cervicogenic headache: an assessment of the evidence on clinical diagnosis, invasive tests, and treatment. The Lancet Neurology. 8: 959-968, 2009.
4. Day M. Facial Pain and Cervicogenic Headache. The 14th Annual Advanced Interventional Pain Conference and Practical Workshop and Interventional Techniques Review Course: 22-23, 2009.

## SANG CHUL LEE, MD, PHD, FIPP

### BIOGRAPHICAL SKETCH

Prof. Sang Chul Lee is a Professor and Chairman of the Department of Anesthesiology and Pain Medicine, Seoul National University College of Medicine, and the President of Korean Spinal Pain Society and Korean IASP chapter.

### ULTRASOUND GUIDED TREATMENT 2011

#### Objectives

Upon completion of this presentation attendees will be able to discuss

- Why we should use ultrasound as a guidance method in pain treatment
- What basic principles of ultrasound imaging are
- For what ultrasound guided is used in the field of pain treatment
- Relationships between the inserted needle and inner structures
- Proper postures during ultrasound guided intervention
- How Sonoanatomy compare with real anatomy
- Examples of ultrasound application for pain treatment

#### Key Points

- Ultrasonography has potential usefulness in pain management including diagnosis and interventional treatment.
- The rationale for performing ultrasound guided treatment is that it provides information that aids in establishing a diagnosis and prognosis, locating areas of pathology, and providing therapy via a real-time visualization.
- Ultrasonography is the only modality that allows direct visualization of relationships between the inserted needle and inner structures such as vessels or nerves in the way of target areas to avoid an iatrogenic injury of them.
- Barriers to the use of ultrasound in clinical practice include necessity of training for operation due to some limitations of ultrasound-guided intervention such as unrecognized intravascular injection.
- Expected outcomes include ruling in or out area or areas of pathology, facilitating treatment, better forecasting of prognosis and future treatment options.

#### References

1. Andres JD, Sala-Blanch X. Ultrasound imaging techniques for regional nerve blocks. In: Interventional Pain Management: Image Guided Procedures, 2nd ed. P Raj et al, eds. Saunders Elsevier, Philadelphia, pp 584-596, 2008.
2. Bianchi S, Martinoli C. Ultrasound of the musculoskeletal system. Springer-Verlag Berlin Heidelberg, New York, 2007.
3. Lee SH et al. Ultrasound guided regional anesthesia & pain intervention. Hansol, Seoul, 2010.
4. Hadzic A. Textbook of regional anesthesia and acute pain management. McGraw-Hill, New York, pp 657-694, 2007.

ADNAN A. AL-KAISEY, MB CHB, FFRCA, FPMRCA, FIPP

### BIOGRAPHICAL SKETCH

Dr Al-Kaisy is currently Clinical Lead and Consultant at the Pain Management and Neuromodulation Centre/ Guy's and St Thomas Hospital. He trained in Chronic Pain Medicine at The Walton Centre, Liverpool for Neurology and Neurosurgery. He has a fellowship in Chronic Pain Management at University of Toronto Hospital, Canada.

He has a number of publications and research in variety of categories in pain management.

His interest is in management of Spine and Neuropathic pain. He has extensive experience in Neuromodulation: Spinal Cord Stimulation for Failed Back Surgery Syndrome, Intractable Angina, Nerve Lesion, and Sacral Nerve Stimulation for Urinary Incontinence, Interstitial Cystitis and Bowel Incontinence. Dr Al-Kaisy was voted the Hospital Doctor of the Year in 2001 for the Pain Management.

### HIGH FREQUENCY SPINAL CORD STIMULATION IN THE MANAGEMENT OF AXIAL BACK PAIN

#### Objectives

Upon completion of this presentation attendees will be able to discuss

- The role of conventional Spinal Cord Stimulation (SCS) in management of Failed back Surgery Syndrome (FBSS).
- Limitations of conventional SCS in the management of Axial Back Pain (ABP)
- Strategies used to improve the efficacy of the conventional SCS
- What is high frequency stimulation?
- How does high frequency stimulation work? How safe is it?
- What are the advantages of high frequency stimulation for patients, operators and providers?
- Future direction of high frequency SCS

#### Key Points

- Spinal Cord Stimulation is evidence-based treatment used in the management of chronic pain conditions.
- While SCS is very effective for radicular pain, one notable area that SCS has had less success in is ABP, which is a mix of nociceptive and neuropathic pain.
- In conventional SCS, paraesthesia coverage has been essential for pain relief. However, coverage of low back pain without dorsal root stimulation and without undesirable stimulation is difficult to accomplish.
- One promising approach for this unmet need is high frequency SCS using up to 10 KHZ.
- In a multi-centre prospective European open label study with 84 implanted patients, high frequency SCS technology showed significant relief for chronic back pain in difficult-to-treat patients, such as predominant back pain patients.
- Leads can be placed in anatomic midline rather than physiologic midline, making the procedure simpler. Paraesthesia mapping step is not required, making the time for high frequency SCS surgery more predictable and potentially shorter.
- Future directions of HF SCS includes use of different algorithms in programming, different applications and advances in equipment technology.

#### References

1. Hill R, Garrett Z, Saile E. Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin. NICE technology appraisal guidance 159. October 2008. <http://www.nice.org.uk/TA159>. National Institute for Health and Clinical Excellence.
2. Kuechmann, et al. Could automatic position adaptive stimulation be useful in spinal cord stimulation. 6th Congress of the European Federation of IASP Chapters 2009 and [www.restoresensor.eu](http://www.restoresensor.eu).
3. Smet I, Van Buyten JP, Al-Kaisy A. European Prospective Study with the Nevro Implantable System. North American Neuromodulation Society 2010 Meeting.

# PNS Therapy for Back Pain: Now CE marked



**BACK  
PAIN**

WHEN IT COMES TO BACK PAIN WE'RE IN FRONT.

## MURRAY H. ROSENTHAL, DO, FAPA

### BIOGRAPHICAL SKETCH

Dr. Murray Rosenthal, a board-certified psychiatrist and Fellow of the American Psychiatric Association, is Chief Medical Officer of Millennium Laboratories. Prior to joining Millennium, Dr. Rosenthal was Senior Medical Director and CEO of California Clinical Trials (CCT), a multi-site, phase 1-4 CNS research group. During his tenure at CCT he conducted over 450 clinical trials in all areas of Central Nervous System (CNS) research.

### THE ROLE OF URINE DRUG TESTING IN PAIN MANAGEMENT

#### Objectives

Upon completion of this presentation attendees will be able to discuss

- The rationale for performing urine drug testing (UDT) on patients taking opioids
- How do Point of Care Urine Testing (POCT) and laboratory confirmation compare
- Limitations of POCT
- What are common objections to UDT testing?
- APS Guidelines for testing; when to test
- How to assess false positives and negatives
- Using UDT to assess you patient and practice; what could unexpected results indicate
- Current and future research

#### Key Points

- Physicians guessing at patient compliance on opioids is less than chance
- UDT is becoming a part of physician-patient treatment contract
- Patients need to understand the UDT as a routine procedure like monitoring Bp in hypertension.
- The POCT and confirmation have different uses based upon their sensitivity. One gives instant snapshot and the other can be used over time to assess levels compare to large patient populations on similar medications.
- Testing should be both routine and at key times during changes in treatment or significant changes in a patient's clinical picture.
- By assessing huge databases of patient samples, we have observed patterns of use and misuse which will be discussed

#### References

1. Michna E, Jamison RN, Pharm-D, et al. Urine toxicology screening among chronic pain patients on opioid therapy, frequency and predictability of abnormal findings. Clin J Pain 2007; 33: 173-175
2. Amadeo Pesce, Ph.D., DABCC, et al. Diagnostic accuracy an interpretation of urine drug testing for pain patients; an evidence-based approach. Toxicology and Drug Testing, Rijeka, Croatia; InTech-Open Access Publisher 2011
3. West R, Pesce A, Mikel C, Rosenthal M, Latyshev S, Crews B and Almazan P. Observations of medication compliance by measurement of urinary drug concentrations in a pain management population. J Opioid Manag 2010;6:253-257.
4. Pesce A, Rosenthal M, West R, West C, Crews B, Mikel C, Almazan P and Latyshev S. An evaluation of the diagnostic accuracy of liquid chromatography-tandem mass spectrometry versus immunoassay drug testing in pain patients. Pain Physician 2010;13:273-281.
5. Pesce A, West C, Rosenthal M, Mikel C, West R, Crews B, Almazan P, Latyshev S and Horn P. Illicit drug use in the pain patient population decreases with more frequent drug testing. Pain Physician 2011;14:189-193.
6. West R, Pesce A, Mikel C, Rosenthal M, Latyshev S, Crews B, Almazan P. Observations of medication compliance by measurement of urine drug concentrations in a pain management population. J Opioid manag 2010; 6: 253-257

## AUTHORS INDEX

<b>ADNAN A. AL-KAISEY</b>	<b>16, 50</b>
<b>RAY M. BAKER</b>	<b>15, 36</b>
<b>HEMMO A. BOSSCHER</b>	<b>15, 40</b>
<b>COSIMO BRUNI</b>	<b>15, 38</b>
<b>SERDAR ERDINE</b>	<b>14, 22</b>
<b>LORAND EROSS</b>	<b>16, 17, 45</b>
<b>JOSEPH D. FORTIN</b>	<b>14, 15, 31, 35</b>
<b>JAMES E. HEAVNER</b>	<b>15, 40, 41</b>
<b>RAFAEL JUSTIZ</b>	<b>14, 27</b>
<b>SANG CHUL LEE</b>	<b>16, 49</b>
<b>LIONG LIEM</b>	<b>16, 48</b>
<b>JOHN NELSON</b>	<b>15, 16</b>
<b>P. PRITHVI RAJ</b>	<b>17</b>
<b>GABOR B. RACZ</b>	<b>14, 15, 16, 41</b>
<b>RICHARD RAUCK</b>	<b>14</b>
<b>OLAV J. J. M. ROHOF</b>	<b>14, 23, 29</b>
<b>MURRAY H. ROSENTHAL</b>	<b>16, 52</b>
<b>RICARDO RUIZ-LÓPEZ</b>	<b>14, 25</b>
<b>ANDREA TRESOT</b>	<b>16, 42</b>
<b>KRIS C. P. VISSERS</b>	<b>15, 16, 33, 43</b>
<b>JAN PETER WARNKE</b>	<b>15, 40</b>

NOTES

NOTES



**CongressLine Ltd.**

H-1065 Budapest, Révay köz 2.

**Contact person:** Sandra Vamos

**Phone:** +361 429 0146

**Fax:** +361 429 0147

**Email:** [vamos@congressline.hu](mailto:vamos@congressline.hu)

**[www.congressline.hu/pain2011](http://www.congressline.hu/pain2011)**  
**[www.worldinstituteofpain.org](http://www.worldinstituteofpain.org)**